

Authors	Title	Journal	Year	Volume	Pages
Lazarus	Thyroid Regulation and Dysfunction in the Pregnant +AFs—Ultrasonographic evaluation of thyroid tissue perfusion in		2000		
Rogula T, Cichon	patients with Graves-Basedow disease as preparation for surgery+AF0-.	Folia Med Cracov	2000	41(3-4)	121-9
Azizi F, Khoshnati M, Bahrainian M, Hedayati	Thyroid function and intellectual development of infants nursed by mothers taking methimazole.	J Clin Endocrinol Metab	2000	85(9)	3233-8
Raber W, Kmen E, Waldhausl W, Vierhapper	Medical therapy of Graves' disease: effect on remission rates of methimazole alone and in combination with triiodothyronine.	Eur J Endocrinol	2000	142(2)	117-24
Akbar DH, Mushtaq MM, Al-Sheik	Etiology and outcome of thyrotoxicosis at a university hospital.	Saudi Med J	2000	21(4)	352-4
Giovanella L, De Palma D, Ceriani L, Vanoli P, Garancini S, Tordiglione M, Calegaro JU, De Freitas Gomes E, Bae SH, Ulyssea R, Casulari	[Radioiodine treatment of hyperthyroidism using a simplified dosimetric approach. Clinical results]. One-year follow-up of Graves' disease treatment by four different protocols of radioiodine administration.	Radiol Med	2000	100(6)	480-3
Casis FC, Perez	Leukocytoclastic vasculitis: a rare manifestation of propylthiouracil allergy.	Endocr Pract	2000	6(4)	329-32
Isotani H, Sanda K, Kameoka K, Takamatsu	McCune-Albright syndrome associated with non-autoimmune type of hyperthyroidism with development of thyrotoxic crisis.	Horm Res	2000	53(5)	256-9
Saenz	Iodine-131 elimination from breast milk: a case report.	J Hum Lact	2000	16(1)	44-6
Ansaldi GL, Pretolesi F, Varaldo E, Meola C, Minuto M, Borgonovo G, Zingrillo M, Torlontano M, Ghiggi MR, Frusciano V, Perez Parras MA, Marin Paton M, Negrillo Cantero AM, Caro Cruz E, Gonzalez Rivera F, Moreno Carazo, Mitsiades N, Poulaki V, Tseleni-Balafouta S, Vestergaard P, Rejnmark L, Weeke J, Mosekilde	Doppler evaluation of intrathyroid arterial resistances during preoperative treatment with Lugol's iodide solution in patients with diffuse toxic goiter. Radioiodine and percutaneous ethanol injection in the treatment of large toxic thyroid nodule: a long-term study.	J Am Coll Surg	2000	191(6)	607-12
Feret BM, Caley	[Amiodarone-induced hyperthyroidism].	An Esp Pediatr	2000	53(4)	377-9
	Fas ligand expression in thyroid follicular cells from patients with thionamide-treated Graves'	Thyroid	2000	10(7)	527-32
	Fracture risk in patients treated for hyperthyroidism.	Thyroid	2000	10(4)	341-8
	Possible hypothyroidism associated with quetiapine.	Ann Pharmacother	2000	34(4)	483-6

Jamieson A, Semple	Successful treatment of Graves disease in pregnancy with Lugol's iodine.	Scott Med J	2000	45(1)	20-1
Claxton S, Sinha SN, Donovan S, Greenaway TM, Hoffman L,	Refractory amiodarone-associated thyrotoxicosis: an indication for thyroidectomy.	Aust N Z J Surg	2000	70(3)	174-8
Zuniga-Gonzalez	[Thyroiditis induced by radioactive iodine (I-131). Report of a case and review of the literature].	Gac Med Mex	2000	136(1)	65-9
Maragliano G, Zuppa AA, Florio MG, Scapillati ME, Girlando P, Crescimbini B, Cafforio C, Noia G,	Efficacy of oral iodide therapy on neonatal hyperthyroidism caused by maternal Graves' disease.	Fetal Diagn Ther	2000	15(2)	122-6
Andrade VA, Gross JL, Maia	The effect of methimazole pretreatment on the efficacy of radioactive iodine therapy in Graves' hyperthyroidism: one-year follow-up of a prospective,	J Clin Endocrinol Metab	2001	86(8)	3488-9;
Akossou SY, Napporn A, Goeh-Akue E, Hillah A, Sokpoh-Diallo K, Soussou B,	[Problems in the management of thyrotoxicosis in Black Africa: the Tongolese experience].	Ann Endocrinol (Paris)	2001	62(6)	516-20
Hao ST, Reasner CA 2nd, Becker	Use of cold iodine in patients with Graves' disease: observations from a clinical practice.	Endocr Pract	2001	7(6)	438-42
Qari FA, Naser TA, Hashim IM, Tevaarwerk	Outcome of thyrotoxicosis treatment with 3 different modalities.	Saudi Med J	2001	22(10)	907-9
Teelucksingh S, Singh V, Balkaran	Use of radioiodine in adolescent Graves' disease.	Ann Trop Paediatr	2001	21(4)	335-8
Cundiff JG, Portugal L, Sarne	Parathyroid adenoma after radioactive iodine therapy for multinodular goiter.	Am J Otolaryngol	2001	22(5)	374-5
Urbannek V, Voth E, Moka D, Schicha	[Radioiodine therapy of Graves' disease--a dosimetric comparison of various therapy regimens of antithyroid agents].	Nuklearmedizin	2001	40(4)	111-5
Nishiyama K, Kitahara A, Natsume H, Matsushita A, Nakano K, Sasaki S, Genma R, Fontanilla JC, Schneider AB, Sarne	Malignant hyperthermia in a patient with Graves' disease during subtotal thyroidectomy.	Endocr J	2001	48(2)	227-32
Clerc J, Kahn E, Fragu	The use of oral radiographic contrast agents in the management of hyperthyroidism. SIMS evidence that carbimazole enhances spatial heterogeneity of thyroid iodine storage and targeting in a woman with Graves'	Thyroid	2001	11(6)	561-7
Sabri O, Zimny M, Schreckenberger M, Reinartz P, Nowak B, Ostwald E, Schafer W,	[Characterization of therapy failures in radioiodine therapy of Graves' disease without simultaneous antithyroid agents].	Cell Mol Biol (Noisy-le-grand)	2001	47(3)	519-27
		Nuklearmedizin	2001	40(1)	1-6

Kadmon PM, Noto RB, Boney CM, Goodwin G, Gruppuso	Thyroid storm in a child following radioactive iodine (RAI) therapy: a consequence of RAI versus withdrawal of antithyroid Side-effects of iodized oil administration in patients with simple goiter.	J Clin Endocrinol Metab	2001	86(5)	1865-7
Azizi F, Daftarian Robert J, Mariethoz S, Pache JC, Bertin D, Caulfield A, Murith N, Peytremann A, Goumaz M, Garcia B, Martin-Du Pan	Short- and long-term results of total vs subtotal thyroidectomies in the surgical treatment of Graves' disease.	J Endocrinol Invest	2001	24(2)	72-7
Baskin HJ, Cobin RH, Duick DS, Gharib H, Guttler RB, Kaplan MM, Segal	American Association of Clinical Endocrinologists medical guidelines for clinical practice for the evaluation and treatment of hyperthyroidism and Thyrotoxicosis—surgical management in the era of evidence-based medicine: experience in western India with 752 cases.	Swiss Surg	2001	7(1)	20-4
Bhansali SK, Chandalia		Endocr Pract	2002	8(6)	457-69
Baskin HJ, Cobin RH, Duick DS, Gharib H, Guttler RB, Kaplan MM, Segal RL, Garber JR, Hamilton CR Jr, Handelsman Y, Hellman R, Kukora JS, Levy P, Palumbo PJ, Petak SM, Rettinger HI, Rodbard HW, Reiners C, Schneider Vidal-Trecan GM, Stahl JE, Durand-Hoermann R, Quadbeck B, Roggenbuck U, Szabolcs I, Pfeilschifter J, Meng W, Reschke K, Hackenberg K, Dettmann J, Prehn B, Hirche H, Mann Bogazzi F, Bartalena L, Campomori A, Brogioni S, Traino C, De Martino F, Rossi G, Lippi F,	AMERICAN ASSOCIATION OF CLINICAL ENDOCRINOLOGISTS MEDICAL GUIDELINES FOR CLINICAL PRACTICE FOR THE EVALUATION AND TREATMENT OF HYPERTHYROIDISM AND HYPOTHYROIDISM.	Asian J Surg	2002	25(4)	291-9
	Radioiodine therapy of thyroid autonomy.	Endocr Pract	2002	8(6)	457-46!
	Managing toxic thyroid adenoma: a cost-effectiveness analysis.	Eur J Nucl Med Mol Imaging	2002	29 Supp	S471-8
		Eur J Endocrinol	2002	146(3)	283-94
	Relapse of Graves' disease after successful outcome of antithyroid drug therapy: results of a prospective randomized study on the use of levothyroxine.	Thyroid	2002	12(12)	1119-21
	Treatment with lithium prevents serum thyroid hormone increase after thionamide withdrawal and radioiodine therapy in patients with Graves' disease.	J Clin Endocrinol Metab	2002	87(10)	4490-5

Tepmongkol	Enhancement of radioiodine uptake in hyperthyroidism with hydrochlorothiazide: a prospective randomised control	Eur J Nucl Med Mol Imaging	2002	29(10)	1307–11
Cardoso LC, Martins DC, Campos DV, Santos LM, Correa da Costa VM, Rosenthal D, Xiao H, Zhuang W, Wang S, Yu B, Chen G, Zhou M, Braga M, Walpert N, Burch HB, Solomon BL, Cooper	Effect of iodine or iopanoic acid on thyroid Ca ²⁺ /NADPA-dependent H ₂ O ₂ -generating activity and thyroperoxidase in toxic diffuse goiters.	Eur J Endocrinol	2002	147(3)	293–8
Yamashita Y, Yamane K, Fujikawa R, Okubo M, Kohno	Arterial embolization: a novel approach to thyroid ablative therapy for Graves' disease.	J Clin Endocrinol Metab	2002	87(8)	3583–9
Thomopoulos	The effect of methimazole on cure rates after radioiodine treatment for Graves' hyperthyroidism: a randomized clinical trial.	Thyroid	2002	12(2)	135–9
Bolanos F, Gonzalez-Ortiz M, Duron H, Sanchez Zumo LA, Terzian C, Brannan Chao CS, Lin SY, Sheu	A successful pregnancy and delivery case of Graves' disease with myeloperoxidase antineutrophil cytoplasmic antibody induced by [Hyperthyroidism due to excess iodine].	Endocr J	2002	49(5)	555–9
Tanwani LK, Lohano V, Broadstone VL, Bogazzi F, Aghini- Lombardi F, Cosci C, Lupi I, Santini F, Tanda ML, Miccoli P, Basolo F, Pinchera A, Bartalena L, Osman F, Franklyn JA, Sheppard MC, Gammie	Remission of Graves' hyperthyroidism treated with methimazole.	Presse Med	2002	31(35)	1664–9
Alexander EK, Larsen	Thyrotoxic hypokalemic periodic paralysis in a Hispanic male.	Rev Invest Clin	2002	54(4)	307–10
Dahlen	Graves' disease presented as painful goiter.	J Natl Med Assoc	2002	94(5)	383–6
Eaton SE, Euinton HA, Newman CM, Weetman AP, Bennet	Minimal change nephropathy and graves' disease: report of a case and review of the literature.	Horm Res	2002	57(1–2)	53–6
Cardenas GA, Cabral JM, Leslie	Iopanoic acid rapidly controls type I amiodarone-induced thyrotoxicosis prior to thyroidectomy.	Endocr Pract	2002	8(1)	40–3
		J Endocrinol Invest	2002	25(2)	176–80
		Circulation	2002	105(11)	1275–7
		J Clin Endocrinol Metab	2002	87(3)	1073–7
		Crit Care Nurse	2002	22(1)	62–9
		Clin Endocrinol (Oxf)	2002	56(1)	33–8
		Cleve Clin J Med	2003	70(7)	624–6, 1

Bojarska-Szmygin A, Janicki K, Pietura R, Janicka	The usefulness of thyroid size and TSH receptor antibody (TRAb) determinations in predicting the effectiveness of tiamazole and I-131 treatment for Graves-Basedow's disease.	Ann Univ Mariae Curie Sklodowska Med	2003	58(1)	242-7
Dai WX, Lian XL, Lu L, Li SM, Li SH, Li	+AFs-Effect of universal salt iodization on the dosage of antithyroid drug+AF0-.	Zhonghua Yi Xue Za Zhi	2003	83(24)	2142-5
Bonnema SJ, Bennedbaek FN, Gram J, Veje A, Marving J, Hegedus	Resumption of methimazole after 131I therapy of hyperthyroid diseases: effect on thyroid function and volume evaluated by a randomized clinical trial.	Eur J Endocrinol	2003	149(6)	485-92
Bogazzi F, Martino E, Dell'Unto E, Brogioni S, Cosci C, Aghini-Lombardi F, Ceccarelli C, Pinchera A,	Thyroid color flow doppler sonography and radioiodine uptake in 55 consecutive patients with amiodarone-induced thyrotoxicosis.	J Endocrinol Invest	2003	26(7)	635-40
Streetman DD, Khanderia	Diagnosis and treatment of Graves disease.	Ann Pharmacother	2003	37(7-8)	1100-9
Erdogan MF, Gulec S, Tutar E, Baskal N, Erdogan	A stepwise approach to the treatment of amiodarone-induced thyrotoxicosis.	Thyroid	2003	13(2)	205-9
Agboola-Abu CF, Kuku	Experience in the use of radioactive iodine therapy for hyperthyroidism in Nigerian patients. A study of twenty-two [Therapeutic specifics in atrial fibrillation in hyperthyroidism. Study of 14 cases].	West Afr J Med	2003	22(4)	324-8
Kamoun M, Haffani M, Basdeh L, Abid H, Kraiem S, Slimane H, Haddad	[Atrial fibrillation in hyperthyroidism. Study of 14 cases].	Tunis Med	2003	81(12)	932-6
Azizi F, Bahrainian M, Khamseh ME, Khoshnati	Intellectual development and thyroid function in children who were breast-fed by thyrotoxic mothers taking methimazole.	J Pediatr Endocrinol Metab	2003	16(9)	1239-4;
Iovicic M, Radiojkovic B, Penezic Z, Stojkovic M, Tancic M, Vujovic	[Agranulocytosis and acute coronary syndrome in apathetic hyperthyroidism].	Srp Arh Celok Lek	2003	131(5-6)	249-53
Bianconcini M, Giannini	[Amiodarone-induced thyrotoxicosis: case report and review of the literature].	Ital Heart J Suppl	2003	4(8)	682-7
Pearce EN, Bogazzi F, Martino E, Brogioni S, Pardini E, Pellegrini G, Parkes AB, Lazarus JH, Kettaneh A, Prevot S, Biaggi A, Stirnemann J, Fain O, Hocqueloux L, Mouas H, Levy VG,	The prevalence of elevated serum C-reactive protein levels in inflammatory and noninflammatory thyroid disease.	Thyroid	2003	13(7)	643-8
	Hyperthyroidism in two patients with Crohn disease and Takayasu arteritis.	Scand J Gastroenterol	2003	38(8)	901-3

Abalovich M, Llesuy S, Gutierrez S, Repetto	Peripheral parameters of oxidative stress in Graves' disease: the effects of methimazole and 131 iodine treatments.	Clin Endocrinol (Oxf)	2003	59(3)	321–7
Aitken M, George A, Bodmer C, Cameron	The efficacy of low-dose radioactive iodine without a thionamide in the treatment of thyrotoxicosis.	Clin Med (Lond)	2003	3(3)	265–7
Ghadban WK, Zirie MA, Al-Khateeb DA, Jayyousi AA, Mobayehd HM, El- Aloosy	Radioiodine treatment of hyperthyroidism. Success rate and influence of thyrostatic medication.	Saudi Med J	2003	24(4)	347–51
Franklyn	Thyrotoxicosis. +AFs—Problems and new developments in the management of Graves' disease: medical therapy+AF0—.	Clin Med (Lond)	2003	3(1)	11–5
Quadbeck B, Janssen OE, Mann	Administration of a single dose of recombinant human thyrotrophin enhances the efficacy of radioiodine treatment of large compressive multinodular goitres.	Z Arztl Fortbild Qualitatssich	2004	98 Supp	37–44
Silva MN, Rubio IG, Romao R, Gebrin EM, Buchpiguel C, Tomimori E, Camargo R, Cardia MS, Medeiros-	Successful treatment of hyperthyroidism with amiodarone in a patient with propylthiouracil-induced acute hepatic failure.	Clin Endocrinol (Oxf)	2004	60(3)	300–8
Brusco F, Gonzalez G, Soto N, Arteaga	Scintigraphy for risk stratification of iodine-induced thyrotoxicosis in patients receiving contrast agent for coronary angiography: a prospective study of patients with low thyrotropin.	Thyroid	2004	14(10)	862–5
Fricke E, Fricke H, Esdorn E, Kammeier A, Lindner O, Kleesiek K, Horstkotte D,	Treating amiodarone-induced thyrotoxicosis with radioactive iodine.	J Clin Endocrinol Metab	2004	89(12)	6092–6
Iskandar SB, Jordan RM, Peiris	[Thyrotoxic crisis in a patient with Graves' disease].	Tenn Med	2004	97(9)	408–10
Prujim MT, Pereira	Therapeutic plasmapheresis in patients with severe hyperthyroidism in whom antithyroid drugs are Iodine repletion, thyrotoxicosis and atrial fibrillation in Isfahan, Iran.	Ned Tijdschr Geneesk	2004	148(34)	1691–4
Ozbey N, Kalayoglu-Besisik S, Gul N, Bozbora A, Sencer E, Aminorroaya A, Rohani S, Sattari G, Haghghi S, Hadj Ali I, Khiari K, Cherif L, Ben Abdallah N, Ben Maiz H, Hajri H, Riggs JE, Pandey HK, Schochet SS Topliss DJ, Eastman	[Treatment of Graves' disease: 300 cases].	Int J Clin Pract	2004	58(6)	554–8
Woenckhaus U, Girlich	Critical illness myopathy associated with hyperthyroidism. 5: Diagnosis and management of hyperthyroidism and [Therapy and prevention of hyperthyroidism].	Ann Saudi Med	2004	24(1)	13–7
		Presse Med	2004	33(1)	17–21
		Mil Med	2004	169(1)	71–2
		Med J Aust	2004	180(4)	186–93
		Internist (Berl)	2005	46(12)	1318–2

Quadbeck B, Hoermann R, Roggenbuck U, Hahn S, Mann K, Janssen	Sensitive thyrotropin and thyrotropin-receptor antibody determinations one month after discontinuation of antithyroid drug treatment as predictors of relapse in Graves' disease.	Thyroid	2005	15(9)	1047-54
Reid JR, Wheeler	Hyperthyroidism: diagnosis and treatment. Usefulness of the 2nd generation assay for anti-TSH receptor antibodies to differentiate relapse of Graves' thyrotoxicosis from development of painless thyroiditis after antithyroid drug treatment for Graves' disease.	Am Fam Physician	2005	72(4)	623-30
Izumi Y, Takeoka K, Amino	Effect of long-term continuous methimazole treatment of hyperthyroidism: comparison with radioiodine.	Endocr J	2005	52(4)	493-7
Azizi F, Ataie L, Hedayati M, Mehrabi Y, Sheikholeslami Minelli R, Valli MA, Di Secli C, Finardi L, Chiadra P, Bertoni R, Ferrari C, Barilli AL, Coiro V, Saccani Jotti G, Kubota S, Ohye H, Sasaki I, Nishihara E, Kudo T, Fukata S, Amino N, Kuma K, Mitsuda N, Miyauchi	Is steroid therapy needed in the treatment of destructive thyrotoxicosis induced by alpha-interferon in chronic hepatitis C?	Eur J Endocrinol	2005	152(5)	695-704
Zantut-Wittmann DE, Ramos CD, Santos AO, Lima MM, Panzan AD, Facuri FV, Etchebehere EC, Lima MC, Kolade VO, Bosinski TJ, Ruffy	Successful use of iodine and levothyroxine to treat Graves' disease in a pregnant patient with allergy to antithyroid drugs and high thyrotropin-binding inhibitor immunoglobulin after radioiodine therapy.	Thyroid	2005	15(12)	1373-6
Rozendaal	High pre-therapy [^{99m} Tc]pertechnetate thyroid uptake, thyroid size and thyrostatic drugs: predictive factors of failure in [¹³¹ I]iodide therapy in Graves' disease.	Nucl Med Commun	2005	26(11)	957-63
Akar S, Comlekci A, Birlik M, Onen F, Sari I, Gurler O, Bekis R, Akkoc Froeschl M, Haddad H, Slot MC, Links TP, Stegeman CA, Tervaert	Acute promyelocytic leukemia after iodine-131 therapy for Graves' disease. [Hyperthyroidism in the elderly: aspecific signs may cause a delay in diagnosis]. Thyrotoxic periodic paralysis in a Turkish male; the recurrence of the attack after radioiodine treatment. Thyrotoxicosis—an uncommon cause of heart failure. Occurrence of antineutrophil cytoplasmic antibodies and associated vasculitis in patients with hyperthyroidism treated with antithyroid drugs: A long-term followup study.	Pharmacotherapy Tijdschr Gerontol Geriatr Endocr J Cardiovasc Pathol Arthritis Rheum	2005 2005 2005 2005 2005	25(7) 36(2) 52(1) 14(1) 53(1)	1017-21 77-80 149-51 24-7 108-13

Nakamura	[Method of medical treatment for Graves' disease based on evidence based medicine].	Nihon Rinsho	2006	64(12)	2243-9
de Souza MV, Buescu A, Vaisman M, de Souza HF, Luiz Dunkelmann S, Kunstner H, Nabavi E, Eberlein U, Groth P, Schumichen Dai WX, Lian XL, Lu L, Li SM, Li SH,	+AFs-The effect of propylthiouracil on the efficacy of radioiodine ($I-131$) therapy in graves hyperthyroidism+AF0-. Lithium as an adjunct to radioiodine therapy in Graves' disease for prolonging the intrathyroidal effective half-life of radioiodine. Useful or not? Effect of universal salt iodization on antithyroid drugs.	Arq Bras Endocrinol Metabol	2006	50(6)	1088-9!
Bonnema SJ, Bennedbaek FN, Veje A, Marving J, Hegedus	Continuous methimazole therapy and its effect on the cure rate of hyperthyroidism using radioactive iodine: an evaluation by a randomized trial.	Nuklearmedizin	2006	45(5)	213-8+
Hiraiwa T, Ito M, Imagawa A, Takamatsu J, Kuma K, Miyauchi A, Hanafusa Ndiaye M, Ndiaye N, Kouamo- Nandjou PN, Deguenonvo R, Ndiaye F, Sy A, Diallo BK, Tall A, Tanrikulu S, Erbil Y, Ademoglu E, Issever H, Barbaros U, Kutluturk F,	Restriction of dietary Iodine does not ameliorate the early effect of anti-thyroid drug therapy for Graves' disease in an area of excessive iodine intake.	Chin Med J (Engl)	2006	119(13)	1108-1:
Cruz Junior AF, Takahashi MH, Albino	[Hyperthyroidism surgery: experience of ENT department of hospital center of Thies].	J Clin Endocrinol Metab	2006	91(8)	2946-5
Fukada	The predictive value of CTLA-4 and Tg polymorphisms in the recurrence of Graves' disease after antithyroid withdrawal.	J Endocrinol Invest	2006	29(4)	380-4
Vijayakumar V, Ali S, Nishino T, Nusynowitz	[Clinical treatment with anti-thyroid drugs or iodine-131 therapy to control the hyperthyroidism of graves disease: a cost-effectiveness analysis].	Dakar Med	2006	51(3)	136-40
Diehl LA, Romaldini JH, Graf H, Bartalena L, Martino E, Albino Vijayakumar V, Nusynowitz ML, Ali	[Toxic multinodular goiter]. What influences early hypothyroidism after radioiodine treatment for Graves' hyperthyroidism?	Endocrine	2006	30(3)	377-81
Fink HJ, Hintze	Management of amiodarone-induced thyrotoxicosis in Latin America: an electronic survey.	Arq Bras Endocrinol Metabol	2006	50(6)	1096-11
Jaruratasirikul S, Leethanaporn K, Sriplung	Is it safe to treat hyperthyroid patients with $I-131$ without fear of thyroid storm? [Current thyroid diagnostics and therapy in disorders of fertility and in pregnancy].	Nihon Rinsho	2006	64(12)	2227-3;
	Thyrotoxicosis in children: treatment and outcome.	Clin Nucl Med	2006	31(11)	688-9
		Clin Endocrinol (Oxf)	2006	65(4)	433-8
		Ann Nucl Med	2006	20(6)	383-5
		Med Klin (Munich)	2006	101(8)	645-52;
		J Med Assoc Thai	2006	89(7)	967-73

Majeed M, Babu	Cholestasis secondary to hyperthyroidism made worse by methimazole.	Am J Med Sci	2006	332(1)	51–3
Patel NN, Abraham P, Buscombe J, Vanderpump	The cost effectiveness of treatment modalities for thyrotoxicosis in a U.K. center.	Thyroid	2006	16(6)	593–8
Mussig K, Thamer C, Bares R, Lipp HP, Haring HU, de Ruijter SH, van Dalen T, Muller	Iodine-induced thyrotoxicosis after ingestion of kelp-containing tea.	J Gen Intern Med	2006	21(6)	C11–4
Tutuncu NB, Tutuncu T, Ozgen A, Erbas	[The surgical treatment of Graves' disease].	Ned Tijdschr Geneeskde	2006	150(24)	1321–5
Bhadada S, Bhansali A, Ohye H, Nishihara E, Sasaki I, Kubota S, Fukata S, Amino N, Kuma K,	Long-term outcome of Graves' disease patients treated in a region with iodine deficiency: relapse rate increases in years	J Natl Med Assoc	2006	98(6)	926–30
Gartner	Juvenile hyperthyroidism: an experience.	Indian Pediatr	2006	43(4)	301–7
Kane MP, Busch	Four cases of Graves' disease which developed after painful Hashimoto's thyroiditis.	Intern Med	2006	45(6)	385–9
Sutradhar SR, Siddiqui NI, Rahman S, Bari MA, Sarker CB, Barca MF, Gruppi C, Oliveira MT, Romao R, Cardia MS, Rubio I, Knobel M, Medeiros-Neto	[Subclinical Hyperthyroidism—What is Expected of the General Physician?].	MMW Fortschr Med	2006	148(9)	30–2
Zhu JQ, Li ZH, Wei T, Zhang H, Gong RX, Xu HZ, Hu LT, Zhang WY, Yang XY, Luo YL, Gong	Drugs for hypothyroidism and hyperthyroidism.	Treat Guidel Med Lett	2006	4(44)	17–24
Zingrillo M, Urbano N, Suriano V, Modoni	Drug-induced thyrotoxic periodic paralysis.	Ann Pharmacother	2006	40(4)	778–81
Dietlein M, Moka D, Reinholtz U, Schmidt M, Schomacker K, Erbil Y, Ozluk Y, Giris M, Salmaslioglu A, Issever H, Barbaros U,	Combined occurrence of hyperthyroid Graves' and Hashimoto's thyroiditis.	Mymensingh Med J	2006	15(1)	102–4
Zhang WY, Yang XY, Luo YL, Gong	Cardiovascular assessment of hyperthyroid patients with multinodular goiter before and after radioiodine treatment preceded by stimulation with recombinant human TSH.	Endocrine	2007	32(2)	175–81
Zhang WY, Yang XY, Luo YL, Gong	+AFs—Study on thyroid defunctionalization method for the preoperative preparation of hyperthyroid operation+AF0–.	Sichuan Da Xue Xue Bao Yi Xue Ban	2007	38(5)	866–70
Zingrillo M, Urbano N, Suriano V, Modoni	Radioiodine treatment of Plummer and multinodular toxic and nontoxic goiter disease by the first approximation dosimetry	Cancer Biother Radiopharm	2007	22(2)	256–60
Dietlein M, Moka D, Reinholtz U, Schmidt M, Schomacker K, Erbil Y, Ozluk Y, Giris M, Salmaslioglu A, Issever H, Barbaros U,	Administration of additional inactive iodide during radioiodine therapy for Graves' disease: who might benefit?	Nuklearmedizin	2007	46(3)	77–84
Zhang WY, Yang XY, Luo YL, Gong	Effect of lugol solution on thyroid gland blood flow and microvessel density in the patients with Graves' disease.	J Clin Endocrinol Metab	2007	92(6)	2182–9

Pedro T, Martinez-Hervas S, Lorente R, Catala M, Ortega J, Real JT, Hiroi N, Sakamoto Y, Urita Y, Higa M, Kuboki K, Yoshino Nishihara E, Nagayama Y, Amino N, Hishinuma A, Takano T, Yoshida Somnuk PH, Pusuwan P, Likitmaskul S, Santiprabhob J, Pacheco Capote C, Mena Bares LM, Benitez Velazco A, Louhibi Rubio L, Contreras Puertas PI, Maza Muret FR, Latre Romero	[Difficult treatment of amiodarone-induced thyrotoxicosis: a case report]. Graves' disease with intractable diarrhea, chylous ascites, and chylothorax: a case report. A novel thyrotropin receptor germline mutation (Asp617Tyr) causing hereditary hyperthyroidism. Treatment outcome of Graves' disease in Thai children. [Usefulness of thyroid scintigraphy in the therapeutic management of amiodarone-induced hyperthyroidism]. [Reflections on mental retardation and congenital hypothyroidism: effects of trace mineral deficiencies]. [Thyroid diseases in sub-Saharan Africa]. Aplasia cutis congenita with skull defect in a monozygotic twin after exposure to methimazole in utero. A new case of familial nonautoimmune hyperthyroidism caused by the M463V mutation in the TSH receptor with anticipation of the disease across generations: a possible role of iodine supplementation. The diagnostic criteria of Graves' disease and especially the thyrotropin receptor antibody; our own experience.	An Med Interna Thyroid Endocr J J Med Assoc Thai Rev Esp Med Nucl Sante Sante Birth Defects Res A Clin Mol Teratol Thyroid Hell J Nucl Med J Pediatr Endocrinol Metab Am J Med Gastroenterol Hepatol	2007 2007 2007 2007 2007 2007 2007 2007 2007 2007 2007 2007 2007 2007 2007 2007 2007 2007 2007	24(11) 17(12) 54(6) 90(9) 26(5) 17(1) 17(1) 79(10) 17(7) 10(2) 20(4) 120(6) 30(5)	543–6 1299–30 927–34 1815–21 270–6 41–50 33–9 680–4 677–80 89–94 535–9 503–5 268–70
Sidibe el	[Neonatal hyperthyroidism: neonatal clinical course of two brothers born to a mother with Graves-Basedow disease, before and after total thyroidectomy.	J Pediatr Endocrinol Metab	2007	20(4)	535–9
Iwayama H, Hosono H, Yamamoto H, Ferrara AM, Capalbo D, Rossi G, Capuano S, Del Prete G, Esposito V, Montesano G, Zampella E, Fenzi G, Salerno M, Paunkovic N, Paunkovic	High-speed heart.	Am J Med	2007	120(6)	503–5
Zuppa AA, Sindico P, Savarese I, D'Andrea V, Fracchiolla A, Cota F, Romagnoli Patel AJ, Collins JS, Patel JR, Casallo Blanco S, Valero MA, Marcos Sanchez F, de Matias Salces L, Blanco Gonzalez JJ, Martin	[Methimazole and propylthiouracil induced acute toxic hepatitis].	Gastroenterol Hepatol	2007	30(5)	268–70

Gietka-Czernel M, Jastrzebska H, Dudek A, Szczepkowski M, Rivkees SA, Dinauer Fukushima T, Tokioka H, Ishii F, Mikane T, Oku S, Fujii H, Katayama D, Kawanishi S, Abe K, Saito H, Takahashi A, Rai T, Takiguchi J, Kanno Y, Monoe K, Young B, Driedger A, Wojcik Sarr A, Ndour MM, Diop SN, Ka CM, Dieng P, Diedhiou D, Sidibe EH, Sow Stalberg P, Svensson A, Hessman O, Nygaard Albino CC, Graf H, Sampaio AP, Vigario A, Paz– El Refaei SM, Shawkat Chemli J, Braham N, Selmi H, Boughattas S, Zbidi A, Essoussi Mazza E, Carlini M, Flecchia D, Blatto A, Zuccarini O, Gamba S, Beninati S, Messina Orsolon P, Giachetti M, Lupi A, Salgarello M, Malfatti V, Zanco Zuberi LM, Awan S, Islam N, Akhter J, Jabbar Malabu UH, Alfadda A, Sulimani RA, Al-Rubeaan KA, Al-Ruhaily AD, Fouda MA, Al– Maatouq MA, El–	[Thyroidectomy as the last chance treatment for life threatening thyrotoxicosis. Case report]. An optimal treatment for pediatric Graves' disease is radioiodine. [Successful anesthetic management of a patient with thyroid storm using landiolol]. [Case of acute hepatitis associated with Basedow's disease]. Autoimmune thyroid disease and segmental myoclonus. [Toxic nodular goiter]. Surgical treatment of Graves' disease: evidence-based approach. Hyperthyroidism (primary). Thiamazole as an adjuvant to radioiodine for volume reduction of multinodular goiter. Long-term carbimazole intake does not affect success rate of radioactive ^{131}I odine in treatment of Graves' hyperthyroidism. [Graves's disease in children and adolescent: study of seven cases]. Long-term follow-up of patients with hyperthyroidism due to Graves' disease treated with methimazole. Comparison of usual treatment schedule with drug discontinuation vs continuous treatment with low methimazole doses: a retrospective study. Pre-therapy hyperfunctioning follicular thyroid carcinoma evaluation with $\text{I}-131$ whole-body scan and with F-18 FDG PET/CT. Does choice of therapy save costs and improve outcomes in hyperthyroid patients? Graves' disease in Saudi Arabia: a ten-year hospital study.	Endokrynol Pol J Clin Endocrinol Metab Masui Nihon Shokakibyo Gakkai Zasshi Mov Disord Dakar Med World J Surg BMJ Clin Evid Expert Opin Investig Drugs Nucl Med Commun Tunis Med J Endocrinol Invest Clin Nucl Med J Pak Med Assoc J Pak Med Assoc	2007 2007 2007 2007 2007 2007 2007 2008 2008 2008 2008 2008 2008 2008 2008 2008 2008 2008 2008	58(1) 92(3) 56(2) 104(1) 22(3) 52(2) 32(7) 2008 17(12) 29(7) 86(8) 31(10) 33(12) 58(6) 58(6)	52–6 797–801 193–5 52–6 434–6 135–40 1269–7 2008 1781–6 642–8 728–34 866–72 882–6 309–12 302–4
--	---	--	--	--	--

Dalan R, Leow MK, Ng	High iodine (substrate) turnover Graves' disease: the intriguing 'rapid responder' variant of	Ann Clin Biochem	2008	45(Pt 6)	612–5
Sanyal D, Mukhopadhyay P, Pandit K, Chatterjee J, Raychaudhuri M, Mukherjee S, Sebastian-Ochoa A, Quesada-Charneco M, Fernandez-Garcia D, Reyes-Garcia R, Rozas-Moreno Mizuno J, Yonenaga K, Mimura Y, Arita H, Hanaoka Watkins MG, Dejhamron P, Huo J, Vazquez DM, Menon	Early treatment with low fixed dose (5 mCi) radioiodine therapy is effective in Indian subjects with Graves' disease.	J Indian Med Assoc	2008	106(6)	360–1, ;
Sanyal P, Bing-You RG, Braverman	Dramatic response to cholestyramine in a patient with Graves' disease resistant to conventional therapy.	Thyroid	2008	18(10)	1115–7
Sato K, Omi Y, Kodama H, Obara T, Yamazaki K, Yamada E, Seki T, Takano K, Shiga T, Pasimeni G, Caroli F, Spriano G, Antonini M, Baldelli R, Appeteccchia Mikosch	[Hyperthermia and metabolic acidosis during subtotal thyroidectomy for a patient with Basedow's disease]. Persistent neonatal thyrotoxicosis in a neonate secondary to a rare thyroid-stimulating hormone receptor activating mutation: case report and literature review. Use of methotrexate to treat isolated Graves ophthalmopathy developing years after thyroidectomy and iodine 131 treatment of papillary thyroid cancer.	Masui	2008	57(7)	897–901
Akin F, Yaylali GF, Bastemir M, Yapar Metso S, Auvinen A, Salmi J, Huhtala H, Jaatinen Galofre Ferrater JC, Corrales Hernandez JJ, Perez Corral B, Canton Blanco A, Alonso Pedrol N, Perez Perez A, Hegedus	Differential diagnosis and appropriate treatment of four thyrotoxic patients with Graves' disease required to take amiodarone due to life- Refractory thyrotoxicosis induced by iodinated contrast agents treated with therapeutic plasma exchange. A case report. [Thyroid and treatment with amiodarone diagnosis, therapy and clinical management]. Effect of methimazole on warfarin anticoagulation in a case of Graves' disease. Increased long-term cardiovascular morbidity among patients treated with radioactive iodine for hyperthyroidism. [Clinical guideline for the diagnosis and treatment of subclinical thyroid dysfunction in pregnancy. Working Group on Subclinical Thyroid Dysfunction of the Spanish Endocrinology Society]. Treatment of Graves' hyperthyroidism: evidence-based and emerging modalities.	Intern Med	2008	47(8)	757–62
		J Clin Apher	2008	23(2)	92–5
		Wien Med Wochenschr	2008	158(1-2)	29–35
		Blood Coagul Fibrinolysis	2008	19(1)	89–91
		Clin Endocrinol (Oxf)	2008	68(3)	450–7
		Endocrinol Nutr	2009	56(2)	85–91
		Endocrinol Metab Clin North Am	2009	38(2)	355–71,

Traisk F, Tallstedt L, Abraham-				
Nordling M,	Thyroid-associated ophthalmopathy after treatment for Graves' hyperthyroidism with antithyroid drugs or iodine-131.	J Clin Endocrinol Metab	2009	94(10) 3700-7
Andersson T, Berg G, Calissendorff J, Hallengren B,				
Hedner P, Lantz M, Nystrom E,				
Ponjavic V, Taube Toni M, Anda E,	[The effects of amiodarone on the thyroid].	An Sist Sanit Navar	2009	32(3) 363-70
Pineda J, Martinez de Esteban JP,				
Guo J, Chen C, Li	[Experiment research of Jiajian Yunjian granules on hyperthyroidism graves].	Zhongguo Zhong Yao Za Zhi	2009	34(18) 2369-7:
Jabrocka-Hybel A, Golkowski F, Baldys-Waligorska A, Sokolowski G, Hubalewska-	Graves' disease with an autonomously functioning thyroid nodule: case report.	Pol Arch Med Wewn	2009	119(7-8) 505-8
Romao R, Rubio IG, Tomimori EK, Camargo RY, Knobel M, Medeiros-Neto	High prevalence of side effects after recombinant human thyrotropin-stimulated radioiodine treatment with 30 mCi in patients with multinodular goiter and subclinical/clinical	Thyroid	2009	19(9) 945-51
Czarnywojtek A, Czepczynski R, Ruchala M, Wasko R, Zgorzalewicz-Stachowiak M, Szczepanek E, Zamyslowska H, Bartkowiak Z, Florek E, Sowinski Yau JS, Chu KS, Li	Radioiodine therapy in patients with amiodarone-induced thyrotoxicosis (AIT).	Neuro Endocrinol Lett	2009	30(2) 209-14
JK, Chan KW, Lau IT, Yum SW, Chan CW, Mo LK, Kwan In H, Pearce EN, Wong AK, Burgess JF, McAneny DB,	Usage of a fixed dose of radioactive iodine for the treatment of hyperthyroidism: one-year outcome in a regional	Hong Kong Med J	2009	15(4) 267-73
Seo JH, Yu HG, Chung	Treatment options for Graves disease: a cost-effectiveness analysis.	J Am Coll Surg	2009	209(2) 170-17!
Ezer A, Caliskan K, Parlakgumus A, Belli S, Kozanoglu Cerqueira C, Knudsen N, Ovesen L, Perrild H, Rasmussen LB, McCormack S, Mitchell DM, Woo M, Levitsky LL, Ross DS, Misra	A case of Vogt-Koyanagi-Harada disease in a patient with Graves disease.	Korean J Ophthalmol	2009	23(2) 112-3
	Preoperative therapeutic plasma exchange in patients with thyrotoxicosis.	J Clin Apher	2009	24(3) 111-4
	Association of iodine fortification with incident use of antithyroid medication--a Danish Nationwide Study.	J Clin Endocrinol Metab	2009	94(7) 2400-5
	Radioactive iodine for hyperthyroidism in children and adolescents: referral rate and response to treatment.	Clin Endocrinol (Oxf)	2009	71(6) 884-91

Yabuta T, Ito Y, Hirokawa M, Fukushima M, Inoue H, Tomoda C, Higashiyama T, Kihara M, Uruno T, Takamura Y, Vejbjerg P, Knudsen N, Perrild H, Laurberg P, Carle A, Pedersen IB, Rasmussen LB, Leow MK, Loh KC, Zhu M, Chan SP, Sundram	Preoperative administration of excess iodide increases thyroid volume of patients with Graves' disease.	Endocr J	2009	56(3)	371–5
		Clin Endocrinol (Oxf)	2009	71(3)	440–5
		Exp Clin Endocrinol Diabetes	2009	117(10)	616–21
Vita R, Lapa D, Vita G, Trimarchi F, Benvenga	A patient with stress-related onset and exacerbations of Graves disease.	Nat Clin Pract Endocrinol Metab	2009	5(1)	55–61
Earl R, Crowther CA, Middleton	Interventions for preventing and treating hyperthyroidism in pregnancy.	Cochrane Database Syst Rev	2010	(9)	CD0086
Nygaard El-Kaissi S, Bowden J, Henry MJ, Yeo M, Champion BL, Oszukowska L, Knapska– Kucharska M, Makarewicz J, Lewinski	Hyperthyroidism (primary). Association between radioiodine therapy for Graves' hyperthyroidism and thyroid-associated ophthalmopathy. The influence of thiamazole, lithium carbonate, or prednisone administration on the efficacy of radioiodine treatment ((131)I) in hyperthyroid patients.	BMJ Clin Evid	2010	2010	
Takata K, Amino N, Kubota S, Sasaki I, Nishihara E, Kudo T, Ito M, Fukata S, Miyauchi	Benefit of short-term iodide supplementation to antithyroid drug treatment of thyrotoxicosis due to Graves' disease.	Int Ophthalmol	2010	30(4)	397–40!
Iwanczuk	[Myocardial infarction and shock associated with thyrotoxicosis].	Endokrynol Pol	2010	61(1)	56–61
Jukic T, Stanicic J, Petric V, Kusic	[Radioiodine versus surgery in the treatment of Graves' hyperthyroidism].	Anestezjol Intens Ter	2010	42(3)	142–6
Sauer-Schulz A, Mussig K, Kurth R, Fend F, Teichmann R, Schnauder	[A rare cause of thyrotoxicosis].	Lijec Vjesn	2010	132(11–)	355–60
Bish LT, Bavaria JE, Augoustides	Thyroid storm after coronary artery bypass grafting.	Dtsch Med Wochenschr	2010	135(37)	1798
Al Jaber J, Haque S, Noor H, Ibrahim B, Al Suwaidi	Thyrotoxicosis and coronary artery spasm: case report and review of the literature.	J Thorac Cardiovasc Surg	2010	140(5)	e67–9
Fumarola A, Di Fiore A, Dainelli M, Grani G, Calvanese	Medical treatment of hyperthyroidism: state of the art.	Angiology	2010	61(8)	807–12
		Exp Clin Endocrinol Diabetes	2010	118(10)	678–84

Boubaker K, Ounissi M, Khiari K, Harzallah A, Abbes M, El Younsi F, Ben Abdellah T, Ben Maiz H, Kinoshita H, Yasuda M, Furumoto Y, Watanabe N, Horiuchi T, Murayama M, Kitamura M, Kaneko S, Inoshita	[A rare case of Graves' disease in a patient on regular haemodialysis].	Nephrol Ther	2010	6(4)	251–4
Chen YT, Yang GG, Hsu	Severe duodenal hemorrhage induced by Lugol's solution administered for thyroid crisis treatment.	Intern Med	2010	49(8)	759–61
Koyuncu A, Aydin C, Topcu O, Gokce ON, Elagoz S, Bogazzi F, Giovannetti C, Fessehatssion R, Tanda ML, Campomori A, Compri E, Rossi G, Ceccarelli C, Vitti	Thyroid storm and lymphocytic myocarditis. Could total thyroidectomy become the standard treatment for Graves' disease? Impact of lithium on efficacy of radioactive iodine therapy for Graves' disease: a cohort study on cure rate, time to cure, and frequency of increased serum thyroxine after antithyroid drug withdrawal. Radioactive iodine-associated changes in thyroid on fine-needle aspiration.	Intern Med	2010	49(6)	593–6
Kapur U, Katz		Surg Today	2010	40(1)	22–5
Bahn RS, Burch HB, Cooper DS, Garber JR, Greenlee MC, Klein I, Laurberg P, McDougall IR, Montori VM, Bahn Chair RS, Burch HB, Cooper DS, Garber JR, Greenlee MC, Klein I, Laurberg P, McDougall IR, Montori VM, Yaqub A, Choudhry MI,	Hyperthyroidism and other causes of thyrotoxicosis: management guidelines of the American Thyroid Association and American Association of Clinical Endocrinologists.	J Clin Endocrinol Metab	2010	95(1)	201–8
Bartalena	Hyperthyroidism and other causes of thyrotoxicosis: management guidelines of the American Thyroid Association and American Association of Clinical Endocrinologists.	Endocr Pract	2011	17(3)	456–521
Pirnat E, Zaletel K, Gaberscek S, Hojker	The outcome of 131I treatment in Graves' patients pretreated or not with methimazole.	Thyroid	2011	21(6)	593–641
Osada E, Hiroi N, Sue M, Masai N, Iga R, Shigemitsu R, Oka R, Miyagi M, Iso K, Kuboki K,	Post-ablative hypothyroidism. The dilemma of how to manage Graves' hyperthyroidism in patients with associated	W V Med J	2011	107(2)	37–40
		J Clin Endocrinol Metab	2011	96(3)	592–9
		Hell J Nucl Med	2011	14(1)	25–9
		Thyroid Res	2011	4(1)	8

Leite AC, Pedro AB, Romaldini	Influence of methimazole and radioactive iodine treatment in the serum levels of the chemokine CXCL10 in hyperthyroid patients with Graves' disease.	Horm Metab Res	2011	43(3)	194–9
Ortiz GA, Lagari-Libhaber V, Prieto-Sanchez LM, Rabinstein	Propylthiouracil-induced lupus, antiphospholipid syndrome, and stroke in a patient with Graves hyperthyroidism.	Arch Neurol	2011	68(12)	1587–91
Orgiazzi	[The treatment of Graves' disease: current views and	Presse Med	2011	40(12 Pt 1)	1155–61
Hou X, Li Y, Li J, Wang W, Fan C, Wang H, Zhang H, Shan Z, Teng	Development of thyroid dysfunction and autoantibodies in Graves' multiplex families: an eight-year follow-up study in Chinese Han pedigrees.	Thyroid	2011	21(12)	1353–8
Noh KW, Seon CS, Choi JW, Cho YB, Park JY, Kim	Thyroid storm and reversible thyrotoxic cardiomyopathy after ingestion of seafood stew thought to contain marine neurotoxin.	Thyroid	2011	21(6)	679–82
Boj Carceller D, Lievano Segundo P, Navarro Beltran P, Sanz Paris A, de Castro Hernandez P, Montreal Villanueva	[Short-term results of treatment with ^{131}I in patients with multinodular goiter: effect of the associated degree of hyperthyroidism and other variables].	Rev Esp Med Nucl	2011	30(3)	156–61
Yang SP, Wu PH, Tey BH, Tan	A patient with thyroid storm presenting with apathetic thyrotoxicosis and features of meningoencephalitis.	Thyroid	2011	21(6)	675–8
Singer J, Eszlinger M, Wicht J, Paschke	Evidence for a more pronounced effect of genetic predisposition than environmental factors on goitrogenesis by a case control study in an area with low normal iodine supply.	Horm Metab Res	2011	43(5)	349–54
Liu JM, Gu LQ, Zhao L, Tang ZY, Sun LH, Hong J, Wang WQ, Luo BY, Zhao YJ, Xu MY, Chen X, Jiang XF, Zhu CM, Jin XL,	Two unusual cases of intractable hyperthyroidism responsive to octreotide: Munchausen syndrome or not?	Clin Chim Acta	2011	412(11–12)	1155–61
Lee HS, Hwang	Seizure and encephalopathy associated with thyroid storm in children.	J Child Neurol	2011	26(4)	526–8
Stellingwerf M, Jellema WT, Eland IA, Wakelkamp	[Agranulocytosis/granulocytopenia after long-term use of thiamazole].	Ned Tijdschr Geneesk	2011	155	A2430
Burch HB, Burman KD, Cooper	A 2011 survey of clinical practice patterns in the management of Graves' disease.	J Clin Endocrinol Metab	2012	97(12)	4549–58
Vila L, Velasco I, Gonzalez S, Morales F, Sanchez E, Lailla JM, Martinez	[Detection of thyroid dysfunction in pregnant women: universal screening is justified].	Med Clin (Barc)	2012	139(11)	509.e1–e6

Santos RB, Romaldini JH, Ward	A randomized controlled trial to evaluate the effectiveness of 2 regimens of fixed iodine ((1)(3)(1)I) doses for Graves disease Short-term effects of combined treatment with potassium bromide and methimazole in patients with Graves' disease.	Clin Nucl Med	2012	37(3)	241–4
Li D, Pei H, Li X, Liu X, Li X, Xie		J Endocrinol Invest	2012	35(11)	971–4
Damle N, Bal C, Kumar P, Reddy R, Virkar	The predictive role of 24h RAIU with respect to the outcome of low fixed dose radioiodine therapy in patients with diffuse toxic	Hormones (Athens)	2012	11(4)	451–7
Campbell K, Doogue	Evaluating and managing patients with thyrotoxicosis.	Aust Fam Physician	2012	41(8)	564–72
Cahn A, Chairsky– Segal I, Olshtain– Pops K, Maayan S, Wolf D, Dresner– Pollak	Changes in thyroid function in Ethiopian and non–Ethiopian Israeli patients with human immunodeficiency virus infection or acquired immunodeficiency syndrome.	Endocr Pract	2012	18(6)	882–6
Sztal-Mazer S, Nakatani VY, Bortolini LG, Boguszewski CL, Spremovic– Radjenovic S, Gudovic A, Lazovic G, Isgoren S, Dagliz Gorur G, Demir H, Berk	Evidence for higher success rates and successful treatment earlier in Graves' disease with higher radioactive iodine doses.	Thyroid	2012	22(10)	991–5
Czarnywojtek A, Waligorska– Stachura J, Szczepanek E, Zgorzalewicz– Stachowiak M, Bereszynska I, Kurdybacha P, Stangierski A, Harasymczuk J, Moura–Neto A, Mosci C, Santos AO, Amorim BJ, de Lima MC, Etchebehere EC, Tambascia MA, Yoshihara A, Noh J, Yamaguchi T, Ohye H, Sato S, Sekiya K, Kosuga Y, Suzuki M, Matsumoto M, Kunii Y, Watanabe Sato S, Endo K, Iizaka T, Saiki R, Iwaku K, Sato S, Takahashi Y,	Fetal free thyroxine concentrations in pregnant women with autoimmune thyroid disease.	J Clin Endocrinol Metab	2012	97(11)	4014–2
	Radioiodine therapy in Graves' disease: is it possible to predict outcome before therapy?	Nucl Med Commun	2012	33(8)	859–63
	A rare case of Interferon–alpha–Induced Hyperthyroidism in patients with a chronic hepatitis C with granulocytopenia and transaminasemia treated successfully with radioiodine.	C Neuro Endocrinol Lett	2012	33(3)	268–72
	Predictive factors of failure in a fixed 15 mCi 131I-iodide therapy for Graves' disease.	Clin Nucl Med	2012	37(6)	550–4
	Treatment of graves' disease with antithyroid drugs in the first trimester of pregnancy and the prevalence of congenital malformation.	J Clin Endocrinol Metab	2012	97(7)	2396–41
	A case of painless thyroiditis in a very early stage of pregnancy.	Intern Med	2012	51(5)	475–7

Osaki Y, Sakurai K, Arihara Z, Hata M, Fukazawa	Prediction of late (24-hour) radioactive iodine uptake using early (3-hour) uptake values in Japanese patients with Graves' Changing trend in surgical	Endocr J	2012	59(2)	173–7
Yip J, Lang BH, Lo	indication and management for Graves' disease.	Am J Surg	2012	203(2)	162–7
Earl R, Crowther CA, Middleton	Interventions for hyperthyroidism pre-pregnancy and during pregnancy.	Cochrane Database Syst Rev	2013	(11)	CD0086
Shinall MC Jr, Broome JT, Nookala R, Shinall JB, Kiernan C, Nilsson H, Hallengren B, Lantz	Total thyroidectomy for Graves' disease: compliance with American Thyroid Association guidelines may not always be [Hyperthyroidism induced by iodine contrast media is hard to treat. Can lead to delayed therapy of tumors, as shown in two	Surgery	2013	154(5)	1009–1!
Sundares V, Brito JP, Wang Z, Prokop LJ, Stan MN, Murad MH, Maia AL, Scheffel RS, Meyer EL, Mazeto GM, Carvalho GA, Graf H, Vaisman M, Maciel LM, Ramos Watt T, Cramon P, Bjorner JB, Bonnema SJ, Feldt-Rasmussen U, Gluud C, Gram J, Hansen JL, Hegedus L, Knudsen N, Bach- Mortensen P, Nolsoe R, Nygaard Czarnywojtek A, Wolinski K, Wasko R, Florek E, Zgorzalewicz- Stachowiak M, Fichna M, Greberska W, Guzik P, Lodyga M, Mizunoya K, Maruyama T, Fujii T, Nasu S, Tanaka N, Amenomori H, Hashimoto T, Regelmann MO, Sullivan CK,	Comparative effectiveness of therapies for Graves' hyperthyroidism: a systematic review and network meta-analysis. The Brazilian consensus for the diagnosis and treatment of hyperthyroidism: recommendations by the Thyroid Department of the Brazilian Society of Endocrinology and Selenium supplementation for patients with Graves' hyperthyroidism (the GRASS trial): Trials study protocol for a randomized controlled trial.	J Clin Endocrinol Metab	2013	98(9)	3671–7
	Amiodarone-induced thyrotoxicosis in a case of Eisenmenger's syndrome.	Arq Bras Endocrinol Metabol	2013	57(3)	205–32
	[Anesthetic and perioperative management of a patient with uncontrolled thyrotoxicosis undergoing coronary artery bypass grafting surgery].	Neuro Endocrinol Lett	2013	34(8)	767–72
	Thyroid "vise" in an infant with neonatal Graves' disease.	Masui	2013	62(10)	1214–7
		Pediatrics	2013	132(4)	e1048–!

Matsumoto T, Miyakoshi K, Saisho Y, Ishii T, Ikenoue S, Kasuga Y, Kadohira I, Sato S, Momotani N, Minegishi K, Catania A, Guaitoli E, Carbotta G, Bianchini M, Di Matteo FM, Carbotta S, Nardi M, Fabiani E, Grani Shinall MC Jr, Broome JT, Baker A, Solorzano Namwongprom S, Unachak K, Dejkhamron P, Ua- apisitwong S, Ekmahachai	Antenatal management of recurrent fetal goitrous hyperthyroidism associated with fetal cardiac failure in a pregnant woman with persistent high levels of thyroid-stimulating hormone receptor antibody after ablative	Endocr J	2013	60(12)	1281–7
Hjorringgaard J, Birkelund T, Ravn	Total thyroidectomy for Graves' disease treatment.	Clin Ter	2013	164(3)	193–6
Hoang TD, Mai VQ, Clyde PW, Shakir Azevedo FV, Blotta FG, Goirgetta JM, Noe R, Vaisman Tanda ML, Piantanida E, Liparulo L, Veronesi G, Lai A, Sassi L, Pariani N, Gallo D, Azzolini C, Lim KK, Wong M, Mohamud WN, Kamaruddin Cascino MD, McNabb B, Gardner DG, Woeber KA, Fox Rakus-Kwiatosz A, Dorosz-Hrenczuk B, Ben-Skowronek Gerber PA, Soll C, Schmid	Is potassium iodide solution necessary before total thyroidectomy for Graves Radioactive iodine for thyrotoxicosis in childhood and adolescence: treatment and outcomes. A patient who experienced thyroid storm complicated by rhabdomyolysis, deep vein thrombosis, and a silent pulmonary embolism: a case [Cardiac arrest and cardiac insufficiency as result of thyrotoxicosis]. Over-the-counter-drug-induced thyroid disorders. [Emerging comorbidities in Graves' disease patients treated with radioiodine with more than 10 years of follow-up]. Prevalence and natural history of Graves' orbitopathy in a large series of patients with newly diagnosed graves' hyperthyroidism seen at a single center. Iodized salt supplementation and its effects on thyroid status amongst Orang Asli in Hulu Selangor, Malaysia. Acute liver failure with thyrotoxicosis treated with liver transplantation. The goiter of a newborn from a mother with Graves disease. Rapidly vanishing hyperthyroidism.	Ann Surg Oncol J Clin Res Pediatr Endocrinol BMC Res Notes Ugeskr Laeger Endocr Pract Arq Bras Endocrinol Metabol J Clin Endocrinol Metab Asia Pac J Clin Nutr Endocr Pract Pediatr Endocrinol Diabetes Metab QJM	2013 2013 2013 2013 2013 2013 2013 2013 2013 2013 2013 2013 2013 2013 2013	20(9) 5(2) 6 175(14) 19(2) 57(1) 98(4) 22(1) 19(3) 19(2) 106(1)	2964–7 95–7 198 962–3 268–74 51–6 1443–9 41–7 e57–60 82–4 69–70

Bianco AC, Anderson G, Forrest D, Galton VA, Gereben B, Kim BW, Kopp PA, Liao XH, Obregon MJ, Peeters RP, Refetoff S, Sharlin	American Thyroid Association Guide to investigating thyroid hormone economy and action in rodent and cell models.	Thyroid	2014	24(1)	88–168
Karyampudi A, Hamide A, Halanaik D, Sahoo JP, Arduc A, Dogan BA, Ozuguz U, Tuna MM, Gokay F, Tutuncu YA, Isik S, Aydin Y, Peksoy	Radioiodine therapy in patients with Graves' disease and the effects of prior carbimazole	Indian J Endocrinol Metab	2014	18(5)	688–93
Moleti M, Violi MA, Montanini D, Trombetta C, Di Bella B, Sturniolo G, Presti S, Alibrandi A, Campenni A,	The effect of radioactive iodine treatment on ¹⁴ C urea breath test results in patients with hyperthyroidism.	Clin Nucl Med	2014	39(12)	1022–6
Mohlin E, Filipsson Nystrom H, Eliasson	Radioiodine ablation of postsurgical thyroid remnants after treatment with recombinant human TSH (rhTSH) in patients with moderate-to-severe graves' orbitopathy (GO): a prospective, randomized, single-blind clinical	J Clin Endocrinol Metab	2014	99(5)	1783–9
Leustean L, Preda C, Ungureanu MC, Danila R, Mogos V, Stefanescu C, Kikuchi S, Ogata T, Fukae J, Tsugawa J, Ouma S, Tanabe M, Yanase T, Tsuboi	Long-term prognosis after medical treatment of Graves' disease in a northern Swedish population 2000–2010.	Eur J Endocrinol	2014	170(3)	419–27
Czarnywojtek A, Warmuz– Stangierska I, Wolinski K, Plazinska M, Kobylecka M, Kunikowska J, Stangierski A, Miechowicz I, Waligorska–	Jod-Basedow effect due to prolonged use of lugol solution— case report.	Rev Med Chir Soc Med Nat Iasi	2014	118(4)	1013–7
Cepkova J, Gabalec F, Svilias I, Horacek	[Prolonged psychosis associated with thyrotoxicosis during the subacute stage of cardioembolic stroke in a patient with untreated basedow disease].	Brain Nerve	2014	66(12)	1509–1·
Cepkova J, Horacek J, Vizda	Radioiodine therapy in patients with type II amiodarone-induced thyrotoxicosis.	Pol Arch Med Wewn	2014	124(12)	695–70·
So E, Arakaki	[The occurrence of agranulocytosis due to antithyroid drugs in a cohort of patients with Graves disease treated with radioactive iodine ¹³¹ I during 14	Vnitr Lek	2014	60(10)	832–6
Weinstock RJ, Lewis T, Miller J,	Radioiodine treatment of Graves' disease – dose/response analysis. Toxic multinodular goiter in a patient with end-stage renal disease and hemodialysis.	Acta Medica (Hradec Kralove)	2014	57(2)	49–55
	Thyroid crisis in the maxillofacial trauma patient.	Hawaii J Med Public Health	2014	73(7)	217–20
		J Oral Maxillofac Surg	2014	72(11)	2148.e1

Schneider DF, Sonnerman PE, Jones MF, Ojomo KA, Chen H, Jaume JC, Elson	Failure of radioactive iodine in the treatment of hyperthyroidism.	Ann Surg Oncol	2014	21(13)	4174–81
El-Kareem MA, Derwish WA, Moustafa	Response rate and factors affecting the outcome of a fixed dose of RAI-131 therapy in Graves' disease: a 10-year study.	Nucl Med Commun	2014	35(9)	900–7
Listernick	A 4-year-old girl with abdominal pain and fever.	Pediatr Ann	2014	43(5)	167–70
Lopez-P Rdel P, Forero JD, Sierra	[Methimazole-induced cholestatic jaundice in a hyperthyroid patient].	Acta Gastroenterol	2014	44(1)	52–8
Liu M, Jing D, Hu J, Yin	Predictive factors of outcomes in personalized radioactive iodine ((131)I) treatment for Graves' disease.	Am J Med Sci	2014	348(4)	288–93
Toderian AB, Lawson	Use of antihistamines after serious allergic reaction to methimazole in pediatric Graves' disease.	Pediatrics	2014	133(5)	e1401–
Albert BB, Eckersley LG, Skinner JR,	QT prolongation in a child with thyroid storm.	BMJ Case Rep	2014	2014	
Matta A, Koppala J, Gossman	Thyrotoxic hypokalaemic periodic paralysis: a rare presentation of Graves' disease in a Hispanic patient.	BMJ Case Rep	2014	2014	
Bisschop PH, van Trotsenburg	Images in clinical medicine. Neonatal thyrotoxicosis.	N Engl J Med	2014	370(13)	1237
Patel N, Inder WJ, Sullivan C, Kaye	An audit of amiodarone-induced thyrotoxicosis—do anti-thyroid drugs alone provide adequate treatment?	Heart Lung Circ	2014	23(6)	549–54
Krol A, Koehler A, Nowak M, Paliczka-Cieslik E, Krajewska J, Kalemba M, Jurecka- Lubieniecka B, Hasse-Lazar K, Michalik B, Szpak- Frankel AJ, Ahmad M, Perl M, Yao J, Torimoto K, Okada Y, Arao T, Mori H, Yamamoto S, Narisawa M, Kurozumi A, Ohye H, Minagawa A, Noh JY, Mukasa K, Kunii Y, Watanabe N, Matsumoto M, Suzuki M, Burch HB, Cooper	Radioactive iodine (RAI) treatment of hyperthyroidism is safe in patients with Graves' orbitopathy—a prospective study.	Endokrynol Pol	2014	65(1)	40–5
	Acute onset of a vesiculopustular rash in an ICU patient.	Dermatol Online J	2014	20(1)	21249
	Glucose variability before and after treatment of a patient with Graves' disease complicated by diabetes mellitus: assessment by continuous glucose monitoring.	Endocr J	2014	61(4)	321–8
	Antithyroid drug treatment for graves' disease in children: a long-term retrospective study at a single institution.	Thyroid	2014	24(2)	200–7
	Management of Graves Disease: A Review.	JAMA	2015	314(23)	2544–5.

Biondi B, Bartalena L, Cooper DS, Hegedus L, Srinivasan S, Misra Schott	The 2015 European Thyroid Association Guidelines on Diagnosis and Treatment of Endogenous Subclinical Hyperthyroidism in children. [Therapy of thyroid nodules].	Eur Thyroid J Pediatr Rev Dtsch Med Wochenschr	2015 4(3) 2015 36(6) 2015 140(8)	149–63 239–48 573–7
Zakavi SR, Khazaei G, Sadeghi R, Ayati N, Davachi B, Bonakdaran S, Jabbari Nooghabi Kyrilli A, Tang BN, Huyge V, Blocklet D, Goldman S, Corvilain B, Sato S, Noh JY, Sato S, Suzuki M, Yasuda S, Matsumoto M, Kunii Y, Mukasa K, Sugino K, Ito K, Nagataki S, Korte AK, Derde L, van Wijk J, Tjan Naqvi SY, Luebbert JJ, Min SH, Phung A, Oh TJ, Han KS, Kim MJ, Kim JM, Lee JH, Park Santarosa VA, Orlandi DM, Fiorin LB, Kasamatsu TS, Furuzawa GK, Kunii IS, Padovani RP, Marone MM, Castiglioni ML, Vieira JG, Maciel RM, Dias-da-Silva Maiga I, Valdes- Socin H, Thiry A, Delwaide J, Sidibe Kaysin A, Viera Yoshihara A, Noh JY, Watanabe N, Mukasa K, Ohye H, Suzuki M, Matsumoto M, Kunii Y, Suzuki N, Kameda T, Iwaku Alswat Siddiqui FA, Sheikh	Methimazole discontinuation before radioiodine therapy in patients with Graves' disease. Thiamazole Pretreatment Lowers the (131)I Activity Needed to Cure Hyperthyroidism in Patients With Nodular Goiter. Comparison of efficacy and adverse effects between methimazole 15 mg+inorganic iodine 38 mg/day and methimazole 30 mg/day as initial therapy for Graves' disease patients with moderate to severe sudden cardiac arrest as a presentation of Brugada syndrome unmasked by thyroid storm. Thyroid storm in a patient with Wolff–Parkinson–White syndrome. Therapeutic Plasmapheresis Enabling Radioactive Iodine Treatment in a Patient with Thyrotoxicosis. Low iodine diet does not improve the efficacy of radioiodine for the treatment of Graves' disease. [Alpha interferon induced hyperthyroidism: a case report and review of the literature]. A case of atypical <i>Bartonella henselae</i> infection in a patient with methimazole-induced agranulocytosis. Substituting Potassium Iodide for Methimazole as the Treatment for Graves' Disease During the First Trimester May Reduce the Incidence of Congenital Anomalies: A Retrospective Study at a Single Medical Institution in Role of Cholestyramine in Refractory Hyperthyroidism: A Case Report and Literature Muscle paralysis in thyrotoxicosis.	Nucl Med Commun J Clin Endocrinol Metab Thyroid BMJ Case Rep BMJ Case Rep J Korean Med Sci Arch Endocrinol Metab Rev Med Liege BMJ Case Rep Thyroid Am J Case Rep BMJ Case Rep	2015 36(12) 2015 100(6) 2015 25(1) 2015 2015 2015 2015 2015 30(10) 2015 59(6) 2015 70(7–8) 2015 25(10) 2015 16 2015 2015	1202–7 2261–7 43–50 1531–4 501–6 390–4 1155–6 486–90

Shin T, Guerrero	Calcium maelstrom: recalcitrant hypocalcaemia following rapid correction of thyrotoxicosis, exacerbated by pregnancy.	BMJ Case Rep	2015	2015
Janson JA, de Laat P, Draaisma	Migratory polyarthritis as an adverse effect of thiamazole use in a 13-year-old girl with Graves'	J Pediatr Endocrinol Metab	2015	28(9–10) 1169–7
Szumowski P, Abdelrazeq S, Kociura Sawicka A, Mojsak M, Kostecki J, Sykala M, Correia MF, Maria AT, Prado S, Limbert, West JD, Cheetham TD, Dane C, Natarajan	Radioiodine therapy for Graves' disease – retrospective analysis of efficacy factors.	Endokrynol Pol	2015	66(2) 126–31
Moraco AH, Banauch G, Kopeć Stein JD, Childers D, Gupta S, Talwar N, Nan B, Lee BJ, Smith TJ, Douglas Liang CM, Ho MH, Wu XY, Hong ZJ, Hsu SD, Chen Hwang S, Shin DY, Song MK, Lee Li C, Tan J, Zhang G, Meng Z, Wang R, Li W, Zheng Uchida N, Suda T, Ishiguro Li HX, Xiang N, Hu WK, Jiao Satoh T, Isozaki O, Suzuki A, Wakino S, Iburi T, Tsuboi K, Kanamoto N, Otani H, Furukawa Y, Teramukai S, Ross DS, Burch HB, Cooper DS, Greenlee MC, Laurberg P, Maia AL, Rivkees SA, Samuels M, Sosa De Leo S, Lee SY, Braverman	Neonatal thyrotoxicosis caused by maternal autoimmune hyperthyroidism. Should radioiodine be the first-line treatment for paediatric Graves' disease? A 46-Year-Old Woman With Dyspnea From an Inhalational Exposure, Triggering Thyroid Storm and Subsequent Multi-Organ System Failure. Risk factors for developing thyroid-associated ophthalmopathy among individuals with Graves disease. Thyroid storm following trauma: a pitfall in the emergency department. High cut-off value of a chimeric TSH receptor (Mc4)-based bioassay may improve prediction of relapse in Graves' disease for Risk factors of hyperthyroidism with hepatic function injury: a 4-year retrospective study. Thyroidectomy in a patient with thyroid storm: report of a case. Relation between therapy options for Graves' disease and the course of Graves' ophthalmopathy: a systematic review and meta-analysis. 2016 Guidelines for the management of thyroid storm from The Japan Thyroid Association and Japan Endocrine Society (First edition). 2016 American Thyroid Association Guidelines for Diagnosis and Management of Hyperthyroidism and Other Causes of Thyrotoxicosis. Hyperthyroidism.	BMJ Case Rep J Pediatr Endocrinol Metab Endokrynol Pol J Pediatr Endocrinol Metab J Intensive Care Med JAMA Ophthalmol Injury Endocrine Horm Metab Res Surg Today J Endocrinol Invest Endocr J Thyroid Lancet	2015 2015 2015 2015 2015 2015 2015 2015 2015 2016 2016	2015 28(7–8) 797–80 30(8) 518–20 133(3) 290–6 46(1) 169–71 48(1) 89–95 47(3) 209–13 45(1) 110–4 39(11) 1225–1 63(12) 1025–11 26(10) 1343–1 388(100) 906–91

Bartalena L, Masiello E, Magri F, Veronesi G, Bianconi E, Zerbini F, Gaiti M, Spreafico E, Gallo D, Premoli P, Piantanida E, Taieb D, Bournaud C, Eberle MC, Catargi B, Schvartz C, Cavarec MB, Faugeron I, Toubert ME, Benisvy D, Archange C,	The phenotype of newly diagnosed Graves' disease in Italy in recent years is milder than in the past: results of a large observational longitudinal study.	J Endocrinol Invest	2016	39(12)	1445–1·
	Quality of life, clinical outcomes and safety of early prophylactic levothyroxine administration in patients with Graves' hyperthyroidism undergoing radioiodine therapy: a randomized controlled study.	Eur J Endocrinol	2016	174(4)	491–50·
Chouhan A, Abhyankar A, Basu	The feasibility of low-dose oral lithium therapy and its effect on thyroidal radioiodine uptake, retention, and hormonal parameters in various subcategories of hyperthyroid 131I therapy for 345 patients with refractory severe hyperthyroidism: Without antithyroid drug pretreatment.	Nucl Med Commun	2016	37(1)	74–8
Ding Y, Xing J, Fang Y, Wang Y, Zhang Y, Long	Hyperthyroidism secondary to hysterosalpingography: an extremely rare complication: A Outpatient 131I Treatment for a Patient with Graves' Disease Receiving Hemodialysis.	Exp Biol Med (Maywood)	2016	241(3)	290–5
Ma G, Mao R, Zhai	No Adverse Effects After Radioiodine Treatment at 3 Weeks of Pregnancy.	Medicine (Baltimore)	2016	95(49)	e5588
Suzuki M, Yoshimura Noh J, Sugino K, Ito Radacic-Aumiler M, Kraljickovic I, Edeljic-Turk V, Makar-Ausperger K, Likic R, Simic Chantra M, Limsuwan A, Mahachoklertwatta Chihara I, Nitta S, Kimura T, Kandori S, Kawahara T, Waku N, Kojima T, Joraku A, Miyazaki J, Iwasaki H, Suzuki H, Kawai K, Leo M, Mautone T, Ionn I, Profilo MA, Sabini E, Menconi F, Mazzi B, Rocchi R, Latrofa F, Nardi M, Vitti P, Marcocci C, Marino	Low cardiac output thyroid storm in a girl with Graves' disease. [Two Cases of Germ Cell Tumors with Hyperthyroidism Due to High Serum hCGLevels]. VARIABLES AFFECTING THE LONG-TERM OUTCOME OF GRAVES ORBITOPATHY FOLLOWING HIGH-DOSE INTRAVENOUS GLUCOCORTICOID PULSE THERAPY IN PATIENTS NOT TREATED WITH ORBITAL	Clin Nucl Med	2016	41(12)	964–96·
		Pediatr Int	2016	58(10)	1080–1·
		Hinyokika Kiyo	2016	62(9)	489–49·
		Endocr Pract	2016	22(10)	1177–1·

Donovan PJ, McLeod DS, Little R, Gordon	Cost–utility analysis comparing radioactive iodine, anti–thyroid drugs and total thyroidectomy for primary treatment of Graves' Interaction of interferon alpha	Eur J Endocrinol	2016	175(6)	595–60;
Gill G, Bajwa H, Strouhal P, Buch	therapy with thyroid function tests in the management of hepatitis C: a case report.	J Med Case Rep	2016	10(1)	253
Xu F, Gu A, Pan Y, Yang L, Ma	Analysis of iodine-131-induced early thyroid hormone variations in Graves' disease.	Nucl Med Commun	2016	37(11)	1154–9
Guan L, Chen G, Zhang J, Wang	The preliminary clinical observation and analysis of childbearingage women with a history of iodine-131 treatment	Biosci Trends	2016	10(4)	307–14
Patel NR, Tamara LA, Lee	99mTc Sestamibi Thyroid Scan in Amiodarone-Induced Thyrotoxicosis Type I.	Clin Nucl Med	2016	41(7)	566–7
Fischli S, Lucchini B, Muller W, Slahor L, Henzen	Rapid preoperative blockage of thyroid hormone production / secretion in patients with Graves' disease.	Swiss Med Wkly	2016	146	w14243
Bruin JE, Saber N, O'Dwyer S, Fox JK, Mojibian M, Arora P, Rezania	Hypothyroidism Impairs Human Stem Cell-Derived Pancreatic Progenitor Cell Maturation in Mice.	Diabetes	2016	65(5)	1297–31
Andersen SL, Olsen J, Laurberg	Maternal thyroid disease in the Danish National Birth Cohort: prevalence and risk factors. RADIONUCLIDE THERAPY WITHOUT RECENT ANTITHYROID DRUG PRETREATMENT FOR HYPERTHYROIDISM COMPLICATED BY SEVERE HYPERBILIRUBINEMIA DUE TO HEPATIC DYSFUNCTION: EXPERIENCE OF A CHINESE	Eur J Endocrinol	2016	174(2)	203–12
Ding Y, Xing J, Qiu Z, Wang Y, Zhang Y, Fang Y, Peng X, Long Y, Deng	Treatment and management of thyroid storm: analysis of the nationwide surveys: The taskforce committee of the Japan Thyroid Association and Japan Endocrine Society for the establishment of diagnostic criteria and nationwide surveys for thyroid storm.	Endocr Pract	2016	22(2)	173–9
Isozaki O, Satoh T, Wakino S, Suzuki A, Iburi T, Tsuboi K, Kanamoto N, Otani H, Furukawa Y, Teramukai S, Akamizu	MANAGEMENT OF ENDOCRINE DISEASE: Subclinical thyrotoxicosis: prevalence, causes and choice of therapy.	Clin Endocrinol (Oxf)	2016	84(6)	912–8
Carle A, Andersen SL, Boelaert K, Laurberg	Comparison of curative effect of 131I and antithyroid drugs in Graves' disease: a meta analysis.	Eur J Endocrinol	2017	176(6)	R325–R
Yuan J, Lu X, Yue	PREDICTION OF RELAPSE FROM HYPERTHYROIDISM FOLLOWING ANTITHYROID MEDICATION WITHDRAWAL USING TECHNETIUM THYROID UPTAKE SCANNING.	Minerva Endocrinol	2017		
Nakhjavani M, Abdollahi S, Farzanefar S, Abousaidi M, Esteghamati A, Naseri M, Eftekhari		Endocr Pract	2017	23(4)	466–471

Chen Y, Liang L, Fang Y, Wang C, Li L, Jiang	[Iodine 131 joint radio frequency ablation treatment for child with hyperthyroidism goiter: one case report].	Zhejiang Da Xue Xue Bao Yi Xue Ban	2017	46(1)	89–91
Yu MG, Quisumbing	Limited cutaneous systemic sclerosis arising in a patient with Graves' disease.	BMJ Case Rep	2017	2017	
Ikeoka T, Otsuka H, Fujita N, Masuda Y, Maeda S, Horie I, Ando T,	Thyroid Storm Precipitated by Diabetic Ketoacidosis and Influenza A: A Case Report and Literature Review.	Intern Med	2017	56(2)	181–18!
Wu VT, Lorenzen AW, Beck AC, Reid VJ, Sugg SL, Howe JR, Pollard JH, Lal G, Weigel	Comparative analysis of radioactive iodine versus thyroidectomy for definitive treatment of Graves disease.	Surgery	2017	161(1)	147–15!

9

8

0

4

1

0

5

ADs- quiz N51-2

2

1

01

2

: quiz 653-4

03

0

372

0

. ix

2

9.e1-2

33

5

0

6

0

2

0

8

.509.e11

1

03

33

5

51

4

3

-7

0

4

4

1

4

233

064

421

8

451

2

313

5

083

3

186

3

09

1337

0

5

5