

Authors	Title	Journal	Year	Volume	Pages
Lin JD, Liou MJ, Chao TC, Weng HF, Ho Seabold JE, Gurl N, Schurrer ME, Aktay R, Kirchner McDougall Franklyn JA, Maisonneuve P, Sheppard M, Betteridge J, Boyle Doi SA, Woodhouse	Prognostic variables of papillary and follicular thyroid carcinoma patients with lymph node Comparison of 99mTc-methoxyisobutyl isonitrile and 201TI scintigraphy for detection Cancer deaths after 131I Cancer incidence and mortality after radioiodine treatment for hyperthyroidism: a population-based cohort study. Ablation of the thyroid remnant and 131I dose in differentiated	Endocr Relat Cancer	1999	6(1)	109–15
Kebebew E, Clark	Medullary thyroid cancer.	J Nucl Med	1999	40(9)	1434–40
Yim JH, Doherty	Papillary thyroid cancer.	Nucl Med	1999	20(5)	407–9
Kebebew E, Clark	Differentiated thyroid cancer: "complete" rational approach.	Lancet	1999	353(917)	2111–5
Griffiths PA, Jones GP, Marshall C, Powley Shingu K, Kobayashi S, Yokoyama S, Fujimori M, Asanuma K, Ito KI, Hama Y, Maruyama M, Kusama	Radiation protection consequences of the care of a terminally ill patient having The likely transformation of papillary thyroid carcinoma into anaplastic carcinoma during postoperative radioactive iodine-131 therapy: report of a	Clin Endocrinol Curr Treat Options	2000	52(6)	765–73
La Quaglia MP, Black T, Holcomb GW 3rd, Sklar C, Azizkhan RG, Haase GM, Newman	Differentiated thyroid cancer: clinical characteristics, treatment, and outcome in patients under 21 years of age who present with distant metastases. A report from the Surgical Discipline Committee	Curr Treat Options World J Surg	2000	1(4)	359–67
Pittas AG, Adler M, Fazzari M, Tickoo S, Rosai J, Larson SM, Robbins Sautter-Bihl ML, Raub J, Hetzel- Sesterheim M, Heinze Chas J, Kowalczyk A, Siekierzynski M, Dziuk E, Janiak Dequanter D, Abdoulaye D, Lothaire P, Gebhart Hadjieva Tutal E, Tutuncu NB, Akcaer N, Bilezikci B, Guvener N, Arican	Bone metastases from thyroid carcinoma: clinical characteristics and prognostic variables in one hundred forty-Differentiated thyroid cancer: prognostic factors and influence of treatment on the [Environmental radiation exposure of a thyroid cancer patient resulting from adjuvant	Br J Radiol Surg Today	2000	73(875)	1209–12
Singer	[Isolated pelvic metastasis of thyroid cancer]. Scoring patients' risk in Unusual case of metastatic thyroid nodule: nonpalpable breast mass as origin. Long-term comparative cancer mortality after use of radio-iodine in the treatment of	J Pediatr Surg	2000	35(6)	910–3
Bal CS, Padhy AK, Kumar	Clinical features of differentiated thyroid carcinoma in children and adolescents	Thyroid	2000	10(3)	955–9; discussion 960
		Strahlenther Onkol	2001	177(3)	261–8
		Wiad Lek Ann Endocrinol (Paris)	2001	54 Suppl	125–31
		Onkologie	2001	62(6)	561–8
		Endocr Pract	2001	7(5)	379–82
		J Insur Med	2001	33(2)	138–42
		Nucl Med Commun	2001	22(8)	881–7

Mukherjee JJ, Kaltsas GA, Islam N, Plowman PN, Foley R, Hikmat J, Britton KE, Jenkins PJ, Chew SL, Monson JP, Besser GM, Al-Balawi IA, Meir HM, Yousef MK, Nayel HA, Al- Thompson LD, Wieneke JA, Paal E, Frommelt RA, Adair CF, Heffess	Treatment of metastatic carcinoid tumours, phaeochromocytoma, paraganglioma and medullary carcinoma of the thyroid with (131)I-meta- iodobenzylguanidine [(131)I- Differentiated thyroid carcinoma referred for radioiodine therapy. A clinicopathologic study of minimally invasive follicular carcinoma of the thyroid gland with a review of the English [The initial treatment of patients with differentiated Differentiated thyroid carcinoma that express sodium-iodide symporter have a lower risk of recurrence for children and adolescents. Childhood and adolescent thyroid carcinoma.	Clin Endocrinol (Oxf)	2001	55(1)	47–60
Smit	[Thyroid papillary microcarcinoma. Salah Azaiz Institute experience].	Saudi Med J	2001	22(6)	497–503
Patel A, Jhiang S, Dogra S, Terrell R, Powers PA, Fenton C, Dinauer CA, Tuttle RM, Francis Grigsby PW, Gal-or A, Michalski JM, Oueslati Z, Aloui M, Gritli S, Touati S, el- May A, Gamoudi A, Ben-Slimene F, Ladgham	Pediatr Res	2001	91(3)	505–24	
Beasley NJ, Lee J, Eski S, Walfish P, Witterick I, Freeman	Ned Tijdschr	2002	146(10)	454–7	
Lee YM, Lo CY, Lam KY, Wan KY, Tam	Cancer	2002	95(4)	724–9	
Gemensjager E, Heitz PU, Martina B, Schweizer Zettinig G, Fueger BJ, Passler C, Kaserer K, Pirich C, Dudczak R, Niederle Chow SM, Law SC, Mendenhall WM, Au SK, Chan PT, Leung TW, Tong CC, Wong IS, Lau	Rev Laryngol Otol Rhinol (Bord)	2002	123(1)	39–42	
Gerard SK, Park Kumar A, Bal	Arch Otolaryngol Head Neck	2002	128(7)	825–8	
Gow KW, Lensing S, Hill DA, Krasin MJ, McCarville MB, Rai SN, Zacher M, Spunt SL, Strickland DK, Eichhorn W, Tabler H, Lippold R, Lochmann M, Schreckenberger M, Bartenstein	J Am Coll Surg	2002	194(6)	711–6	
	Chirurg	2002	73(1)	38–43; discussion 43–!	
	Clin Endocrinol (Oxf)	2002	56(3)	377–82	
	Int J Radiat Oncol Biol Phys	2002	52(3)	784–95	
	J Nucl Med	2003	44(12)	2039–40; author reply	
	Indian J	2003	70(9)	707–13	
	J Pediatr Surg	2003	38(11)	1574–80	
	Thyroid	2003	13(10)	949–58	

Rubino C, de						
Vathaire F, Dottorini						
ME, Hall P, Schwartz	Second primary malignancies in		Br J Cancer	2003	89(9)	1638–44
C, Couette JE,	thyroid cancer patients.					
Dondon MG, Abbas						
MT, Langlois C,						
Reiners	Radioiodine therapy in patients with pulmonary metastases of thyroid cancer: when to treat,	Eur J Nucl Med Mol Imaging	2003	30(7)	939–42	
Besic N, Vidgarn-						
Kralj B, Frkovic-	The role of radioactive iodine in	Thyroid	2003	13(6)	577–84	
Grazio S, Movrin-	the treatment of Hurthle cell carcinoma of the thyroid.					
Stanovnik T,						
Powers PA, Dinauer	Tumor size and extent of disease at diagnosis predict the response to initial therapy for papillary thyroid carcinoma in	J Pediatr Endocrinol Metab	2003	16(5)	693–702	
CA, Tuttle RM, Robie						
DK, McClellan DR,						
Francis						
Gotthardt M, Nowack	Negative correlation between therapeutic success in radioiodine therapy and TcTUs:	Eur J Nucl Med Mol Imaging	2003	30(8)	1165–8	
M, Behe MP,	are TcTUs–adapted dose					
Schipper ML,						
Schlieck A, Hoffken						
Karam M,	Influence of diagnostic and therapeutic doses on thyroid remnant ablation rates.	Nucl Med Commun	2003	24(5)	489–95	
Gianoukakis A,						
Feustel PJ, Cheema						
A, Postal ES, Cooper						
Dorn R, Kopp J, Vogt	Dosimetry–guided radioactive iodine treatment in patients with metastatic differentiated	J Nucl Med	2003	44(3)	451–6	
H, Heidenreich P,						
Carroll RG, Gulec						
Kim TH, Yang DS,	Value of external irradiation for locally advanced papillary thyroid cancer.	Int J Radiat Oncol Biol Phys	2003	55(4)	1006–12	
Jung KY, Kim CY,						
Choi						
Woodings	Radiation protection recommendations for I–131 thyrotoxicosis, thyroid cancer	Australas Phys Eng Sci Med	2004	27(3)	118–28	
Mallick UK,						
Charalambous	Current issues in the management of differentiated	Nucl Med Commun	2004	25(9)	873–81	
Sawka AM,						
Thephamongkhon K,	Clinical review 170: A systematic review and metaanalysis of the effectiveness of radioactive iodine remnant ablation for	J Clin Endocrinol Metab	2004	89(8)	3668–76	
Brouwers M, Thabane						
L, Browman G,						
Gerstein						
Souza Rosario PW,						
Barroso AL, Rezende	Post I–131 therapy scanning in patients with thyroid carcinoma metastases: an unnecessary cost or a relevant contribution?	Clin Nucl Med	2004	29(12)	795–8	
LL, Padrao EL,						
Fagundes TA, Penna						
GC, Purisch						
de Keizer B,						
Hoekstra A,	Bone marrow dosimetry and safety of high 131I activities	J Nucl Med	2004	45(9)	1549–54	
Konijnenberg MW, de	given after recombinant human thyroid-stimulating hormone to treat metastatic differentiated					
Vos F, Lambert B,						
van Rijk PP, Lips CJ,						
Haq MS, McCready	Treatment of advanced differentiated thyroid carcinoma	Nucl Med Commun	2004	25(8)	799–805	
RV, Harmer						
Hu YH, Wang PW,						
Wang ST, Lee CH,						
Chen HY, Chou FF,						
Huang YE, Huang	Influence of 131I diagnostic dose on subsequent ablation in patients with differentiated thyroid carcinoma: discrepancy between the presence of	Nucl Med Commun	2004	25(8)	793–7	

Stokkel MP, Verkooijen RB, Smit Berthe E, Henry- Amar M, Michels JJ, Rame JP, Berthet P, Babin E, Icard P, Samama G, Galateau-Salle F,	Indium-111 octreotide scintigraphy for the detection of non-functioning metastases	Eur J Nucl Med Mol Imaging	2004	31(7)	950-7
	Risk of second primary cancer following differentiated thyroid cancer.	Eur J Nucl Med Mol Imaging	2004	31(5)	685-91
Salvatori M, Perotti G, Rufini V, Maussier ML, Dottorini	Are there disadvantages in administering 131I ablation therapy in patients with differentiated thyroid carcinoma [Poorly differentiated thyroid follicular carcinoma – a clinical Multifactorial analysis on the short-term side effects occurring within 96 hours after radioiodine-131 therapy for [Differentiated thyroid carcinoma in children and adolescents: therapeutic strategy according to clinic Management and outcome of recurrent well-differentiated thyroid carcinoma.	Clin Endocrinol (Oxf)	2004	61(6)	704-10
Liu WS, Qi YF, Tang Kita T, Yokoyama K, Higuchi T, Kinuya S, Taki J, Nakajima K, Michigishi T, Tonami Causeret S, Lifante JC, Borson-Chazot F, Varcus F, Berger N, Peix Palme CE, Waseem Z, Raza SN, Eski S, Walfish P, Freeman Lin JD, Chao TC, Hsueh Chow SM, Law SC, Mendenhall WM, Au SK, Yau S, Mang O, Lau Medvedec	Follicular thyroid carcinomas with lung metastases: a 23-year Differentiated thyroid carcinoma in childhood and adolescence—clinical course and role of radioiodine. Thyroid stunning in vivo and in Is empiric 131I therapy justified for patients with positive thyroglobulin and negative 131I The evolving role of (131)I for the treatment of differentiated	Ai Zheng Ann Nucl Med Ann Chir Arch Otolaryngol Head Neck Endocr J Pediatr Blood Cancer Nucl Med J Nucl Med	2004 2004 2004 2004 2004 2004 2004 2004 2005 2005	23(9) 18(4) 129(6-7) 130(7) 51(2) 42(2) 26(8) 46(7) 46 Suppl	1081-4 345-9 359-64 819-24 219-25 176-83 731-5 1164-70 28S-37S
Ma C, Xie J, Kuang Robbins RJ, Schlumberger Rubino C, Adadj E, Doyon F, Shamsaldin A, Abbas TM, Caillou B, Colonna M, Cecarrelli C, Schwartz C, Bardet S, Langlois C, Ricard M, Schlumberger M, de Vathaire Popova L, Hadjidekova V, Hadjieva T, Agova S, Violot D, M'Kacher R, Adjadj E, Dossou J, de Vathaire F, Pazaitou-Panayiotou K, Kaprara A, Boudina M, Georgiou E, Drimonitis A, Vainas I, Raptou E,	Radiation exposure and familial aggregation of cancers as risk factors for colorectal cancer after radioiodine treatment for thyroid carcinoma. Cytokinesis-block micronucleus test in patients undergoing radioiodine therapy for Evidence of increased chromosomal abnormalities in French Polynesian thyroid Thyroid carcinoma in children and adolescents: presentation, clinical course, and outcome of therapy in 23 children and adolescents in Northern	Int J Radiat Oncol Biol Phys	2005	62(4)	1084-9
		Hell J Nucl Med	2005	8(1)	54-7
		Eur J Nucl Med Mol Imaging	2005	32(2)	174-9
		Hormones (Athens)	2005	4(4)	213-20

Iuchi Y, Sato K, Jimbo J, Inamura J, Shindo M, Ikuta K, Shinzaki H, Ohnishi K, Watanabe S, Podnos YD, Smith D, Wagman LD, Rosa Pelizzo M, Toniato A, Boschin IM, Piotto A, Bernante P, Pagetta C, Palazzi M, Maria Guolo A, Preo P, Williams CE, Woodward Alzahrani AS, Mohamed G, Al Shammary A, Aldasouqi S, Abdal Salam S, Shoukri Leboulleux S, Rubino C, Baudin E, Caillou B, Hartl DM, Bidart JM, Travagli JP, Schlumberger Heymann RS, Brent GA, Hershman Kozak OV, Sukach GG, Korchinskaya OI, Trembach AM, Turicina VL, Voit Rosario PW, Barroso AL, Rezende LL, Padrao EL, Fagundes TA, Reis JS, Purisch Worth AJ, Zuber RM, Hocking Hod N, Hagag P, Baumer M, Sandbank J, Horne Abos Olivares MD, Pesquera Gonzalez Yildirim Panzegrau B, Gordon L, Goudy Rosario PW, Barroso AL, Rezende LL, Padrao EL, Borges MA, Fagundes TA, Rasmuson T, Tavelin Willegaignon J, Stabin MG, Guimaraes MI, Malvestiti LF, Sapienza MT, Maroni	[Acute lymphoblastic leukemia with t(4;11)(q21;q23) after iodine-131 treatment for thyroid cancer]. Radioactive iodine offers survival improvement in Locally advanced differentiated thyroid carcinoma: a 35-year mono-institutional experience in 280 patients. Management of the helpless patient after radioiodine Long-term course and predictive factors of elevated serum thyroglobulin and negative diagnostic radioiodine whole body scan in Prognostic factors for persistent or recurrent disease of papillary thyroid carcinoma with neck lymph node metastases and/or tumor Anaplastic thyroid carcinoma with thyrotoxicosis and Hierarchy of treatment variables affecting outcome of 131I therapy in thyroid cancer patients with lung metastases. Outcome of ablation of thyroid remnants with 100 mCi (3.7 GBq) iodine-131 in patients with thyroid cancer. Radioiodide (131I) therapy for the treatment of canine thyroid Differentiated thyroid carcinoma in children and young adults: evaluation of [Controversies in the follow-up and management of well-differentiated thyroid cancer. A model for predicting outcomes in patients with differentiated thyroid cancer Outpatient therapeutic 131I for thyroid cancer. Ablative treatment of thyroid cancer with high doses of 131I without pre-therapy scanning. Risk of parathyroid adenomas in patients with thyrotoxicosis Evaluation of the potential absorbed doses from patients based on whole-body 131I clearance in thyroid cancer	Rinsho Ketsueki Surgery Nucl Med Commun Nucl Med Commun J Endocrinol Invest J Clin Endocrinol Metab Endocr Pract Exp Oncol Ann Nucl Med Aust Vet J Clin Nucl Med Rev Esp Med Nucl J Am Coll Surg J Nucl Med Technol Nucl Med Commun Acta Oncol Health Phys	2005 2005 2005 2005 2005 2005 2005 2005 2005 2005 2005 2005 2005 2005 2005 2006 2006	46(11) 138(6) 26(11) 26(10) 28(6) 90(10) 11(4) 27(2) 19(3) 83(4) 30(6) 24(3) 200(3) 33(1) 26(2) 45(8) 91(2)	1202-7 1072-6; discussion 10- 965-8 925-8 540-6 5723-9 281-4 150-5 247-50 208-14 387-90 207-15 378-92 28-30 129-32 1059-61 123-7
--	--	---	--	---	--

Jentzen W, Schneider E, Freudenberg L, Eising EG, Gorges R, Muller SP, Brandau W, Bockisch Sioka C, Kouraklis G, Zafirakis A, Manetou A, Dimakopoulos Chatal JF, Campion L, Kraeber-Bodere F, Bardet S, Vuillez JP, Charbonnel B, Rohmer V, Chang CH, Sharkey RM, Goldenberg DM, Kraeber-Bodere F, Rousseau C, Bodet-Milin C, Ferrer L, Faivre-Chauvet A, Campion L, Vuillez JP, Devillers A, Chang CH, Goldenberg DM, Yousuf K, Archibald Grigsby PW, Reddy RM, Moley JF, Hall Chow SM, Yau S, Kwan CK, Poon PC, Law Cappelli C, Pirola I, Braga M, De Martino E, Morassi ML, Gandossi E, Mattanza C, Balzano R, Castellano M, Rosei Verkooijen RB, Smit JW, Romijn JA, Stokkel Hay Robbins RJ, Driedger A, Magner Brzozowska M, Roach Arias F, Vives R, Gomez-Dorronsoro Bhattacharyya N, Chien	Relationship between cumulative radiation dose and salivary gland uptake associated with radioiodine therapy of thyroid cancer. Menstrual cycle disorders after therapy with iodine-131. Survival improvement in patients with medullary thyroid carcinoma who undergo pretargeted anti-carcinoembryonic-antigen radioimmunotherapy: a collaborative study with the Targeting, toxicity, and efficacy of 2-step, pretargeted radioimmunotherapy using a chimeric bispecific antibody and 131I-labeled bivalent hapten in a phase I optimization clinical trial. Brain metastases from papillary adenocarcinoma of the thyroid. Contralateral papillary thyroid cancer at completion thyroidectomy has no impact. Local and regional control in patients with papillary thyroid carcinoma: specific indications of external radiotherapy and Prognostic factors in well-differentiated thyroid carcinoma in patients treated and followed in the same institution. The incidence of second primary tumors in thyroid cancer patients is increased. Selective use of radioactive iodine in the postoperative management of patients with Recombinant human thyrotropin-assisted radioiodine therapy for patients with metastatic thyroid cancer who Timing and potential role of diagnostic I-123 scintigraphy in assessing radioiodine breast uptake before ablation in Cutaneous nodes in a patient with advanced papillary Risk of second primary malignancy after radioactive	Nucl Med Commun Fertil Steril J Clin Oncol J Nucl Med J Otolaryngol Surgery Endocr Relat Cancer Ann Ital Chir Eur J Endocrinol J Surg Oncol Thyroid Clin Nucl Med Clin Transl Oncol Ann Otol Rhinol	2006 27(8) 669-76 2006 86(3) 625-8 2006 24(11) 1705-11 2006 47(2) 247-55 2006 35(6) 366-72 2006 140(6) 1043-7; discussion 10- 2006 13(4) 1159-72 2006 77(2) 107-13 2006 155(6) 801-6 2006 94(8) 692-700 2006 16(11) 1121-30 2006 31(11) 683-7 2006 8(9) 692-3 2006 115(8) 607-10
---	---	---	---

Saghari M, Gholamrezanezhad A, Mirpour S, Eftekhari M, Takavar A, Fard– Esfahani A, Fallahi B, Beiki	Efficacy of radioiodine therapy in the treatment of elevated serum thyroglobulin in patients with differentiated thyroid carcinoma and negative whole– body iodine scan.	Nucl Med Commun	2006	27(7)	567–72
Benbassat CA, Mechlis–Frish S, Hirsch	Clinicopathological characteristics and long–term outcome in patients with Papillary microcarcinoma: is there any difference between	World J Surg	2006	30(6)	1088–95
Lo CY, Chan WF, Lang BH, Lam KY, Segal K, Shpitzer T, Hazan A, Bachar G, Marshak G, Keum KC, Suh YG, Koom WS, Cho JH, Shim SJ, Lee CG, Park CS, Chung WY, Kim	Invasive well–differentiated thyroid carcinoma: effect of treatment modalities on The role of postoperative external–beam radiotherapy in the management of patients with papillary thyroid cancer invading the trachea.	World J Surg	2006	30(5)	759–66
Chuang SC, Hashibe M, Yu GP, Le AD, Cao W, Hurwitz EL, Rao JY, Neugut AI, Walter MA, Turtschi CP, Schindler C, Minnig P, Muller– Brand J, Muller Freudenberg LS, Jentzen W, Marlowe RJ, Koska WW, Luster M, Bockisch Sawka AM, Rotstein L, Brierley JD, Tsang RW, Thabane L, Gafni A, Straus S, Kamalanathan S, Zhao B, Goldstein DP, Rambaldini G, Lin JD, Chao TC, Chen ST, Huang YY, Liou MJ, Hsueh Podnos YD, Smith DD, Wagman LD, Giovannella L, Ceriani L, Ghelfo A, Maffioli M, Keller Saint–Vil D, Emran MA, Lambert R, Alos N, Turpin S, Huot Kucuk NO, Tari P, Tokmak E, Aras Metso S, Auvinen A, Huhtala H, Salmi J, Oksala H, Jaatinen Metso S, Jaatinen P, Huhtala H, Auvinen A, Oksala H, Salmi	Radiotherapy for primary thyroid cancer as a risk factor for second primary cancers. The dental safety profile of high–dose radioiodine therapy for thyroid cancer: long–term results of a longitudinal cohort 124–iodine positron emission tomography/computed tomography dosimetry in pediatric patients with	Int J Radiat Oncol Biol Phys	2006	65(2)	474–80
		Cancer Lett	2006	238(1)	42–52
		J Nucl Med	2007	48(10)	1620–5
		Exp Clin Endocrinol Diabetes	2007	115(10)	690–3
		Thyroid	2007	17(12)	1235–42
		Surg Oncol	2007	16(2)	107–13
		J Surg Oncol	2007	96(1)	3–7
		Clin Endocrinol (Oxf)	2007	67(4)	547–51
		J Pediatr Surg	2007	42(5)	853–6
		Clin Nucl Med	2007	32(4)	279–81
		Cancer	2007	109(10)	1972–9
		J Clin Endocrinol Metab	2007	92(6)	2190–6

Hindie E, Zanotti-Fregonara P, Duron F, Keller I, Bouchard P, Devaux	Should 'low-risk' thyroid cancer patients with residual thyroglobulin be re-treated with iodine 131?	Clin Endocrinol (Oxf)	2007	66(3)	329–34
Reiners C, Demidchik YE, Drozd VM, Biko	Thyroid cancer in infants and adolescents after Chernobyl.	Minerva Endocrinol J Otolaryngol	2008	33(4)	381–95
Page C, Biet A, Zaatar R, Charlet L, Azrif M, Slevin NJ, Sykes AJ, Swindell R, Yap	[Management of the papillary microcarcinoma of the thyroid	2008	37(5)	649–56	
Lin JD, Lin KJ, Chao TC, Hseuh C, Tsang Brown AP, Chen J, Hitchcock YJ, Szabo A, Shrieve DC, Tward Schlumberger MJ, Pacini	Patterns of relapse following radiotherapy for differentiated thyroid cancer: implication for Therapeutic outcomes of papillary thyroid carcinomas	Radiother Oncol	2008	89(1)	105–13
Schneider AB, Viana MA, Ron	The risk of second primary malignancies up to three decades after the treatment of The low utility of pretherapy scans in thyroid cancer	Radiother Oncol J Clin Endocrinol Metab	2008	89(1)	97–104
Seaberg RM, Eski S, Freeman	Weighing shadows: can meta-analysis help define the risk–benefit ratio of RAI treatment Influence of previous radiation exposure on pathologic features and clinical outcome in patients	Thyroid	2009	19(8)	815–6
Sawka AM, Thabane L, Parlea L, Ibrahim-Zada I, Tsang RW, Brierley JD, Straus S, Ezzat S, Goldstein	Second primary malignancy risk after radioactive iodine treatment for thyroid cancer: a systematic review and meta-analysis.	Arch Otolaryngol Head Neck	2009	135(4)	355–9
Riemann B, Schober	Therapeutic strategy of papillary microcarcinoma of the	Thyroid	2009	19(5)	451–7
Biermann M, Pixberg M, Riemann B, Schuck A, Heinecke A, Schmid KW, Willich N, Dralle H, Schober	Clinical outcomes of adjuvant external-beam radiotherapy for differentiated thyroid cancer – results after 874 patient-years of follow-up in the MSDS-trial.	Minerva Endocrinol	2009	34(1)	81–7
Chianelli M, Todino V, Graziano FM, Panunzi C, Pace D, Guglielmi R, Signore A, Papini Duque-Fisher CS, Casiano R, Velez-Hoyos A, Londono-Bustamente	Low-activity (2.0 GBq; 54 mCi) radioiodine post-surgical remnant ablation in thyroid cancer: comparison between [Metastasis to the sinonasal region].	Nuklearmedizin	2009	48(3)	89–98; quiz N15
Maatouk J, Barklow TA, Zakaria W, Al-Abbadia Luster M, Handkiewicz-Junak D, Grossi A, Zacharin M, Taieb D, Cruz O, Hitzel A, Casas JA, Mader U, Dottorini Rios A, Manuel Rodriguez J, Balsalobre MD, Febrero B, Tebar J, Parrilla	Anaplastic thyroid carcinoma arising in long-standing multinodular goiter following Recombinant thyrotropin use in children and adolescents with differentiated thyroid cancer: a multicenter retrospective study. [Distant metastases as the initial manifestation of follicular thyroid carcinoma].	Eur J Endocrinol	2009	160(3)	431–6
		Acta Otorrinolaringol Esp	2009	60(6)	428–31
		Acta Cytol	2009	53(5)	581–3
		J Clin Endocrinol Metab	2009	94(10)	3948–53
		Endocrinol Nutr	2009	56(4)	213–4

Chen PV, Osborne R, Ahn E, Avitia S, Juillard Mihailovic JM, Stefanovic LJ, Malesevic MD, Erak MD, Tesanovic Leenhardt	Adjuvant external-beam radiotherapy in patients with high-risk well-differentiated Metastatic differentiated thyroid carcinoma: clinical management and outcome of disease in patients with initial [Management of thyroid A vision for the surgical management of papillary thyroid carcinoma: extensive lymph node compartmental	Ear Nose Throat J Nucl Med Commun J Radiol J Clin Endocrinol Metab	2009 2009 2009 2009	88(7) 30(7) 90(3 Pt 1) 94(4)	E01 558–64 354–61 1086–8
Arora N, Turbendian HK, Kato MA, Moo TA, Zarnegar R, Fahey TJ 3	Papillary thyroid carcinoma and microcarcinoma: is there a need to distinguish the two?	Thyroid	2009	19(5)	473–7
Gorgone S, Campenni A, Calbo E, Catalfamo A, Sciglitano P, Sofia L, Niceta M, Borzi R, Naing S, Collins BJ, Schneider Miccoli P, Pinchera A, Materazzi G, Biagini A, Berti P, Faviana P, Molinaro E, Viola D, Elisei Lazar L, Lebenthal Y, Steinmetz A, Yackobovitch-Gavan M, Phillip Terezakis SA, Lee KS, Ghossein RA, Rivera M, Tuttle RM, Wolden SL, Zelefsky MJ, Wong RJ, Patel SG, Pfister DG, Mendelsohn AH, Elashoff DA, Abemayor E, St John Sacks W, Fung CH, Chang JT, Waxman A, Braunstein Pawelczak M, David R, Franklin B, Kessler M, Lam L, Shah Panigrahi B, Roman SA, Sosa	[Differentiated thyroid cancers]. G Chir Clinical behavior of radiation- induced thyroid cancer: factors Surgical treatment of low- and intermediate-risk papillary thyroid cancer with minimally invasive video-assisted thyroidectomy. Differentiated thyroid carcinoma in pediatric patients: comparison of presentation and course between pre-pubertal Role of external beam radiotherapy in patients with advanced or recurrent nonanaplastic thyroid cancer: Memorial Sloan-Kettering Cancer Center experience. Surgery for papillary thyroid carcinoma: is lobectomy enough? The effectiveness of radioactive iodine for treatment of low-risk thyroid cancer: a Outcomes of children and adolescents with well- differentiated thyroid carcinoma and pulmonary metastases Medullary thyroid cancer: are practice patterns in the United States discordant from	Thyroid J Clin Endocrinol Metab J Pediatr Int J Radiat Oncol Biol Phys Arch Otolaryngol Head Neck Thyroid Thyroid Ann Surg Oncol	2009 2009 2009 2009 2009 2009 2010 2010 2010 2010 2010 2010 2010	30(1–2) 19(5) 94(5) 154(5) 73(3) 136(11) 20(11) 20(10) 17(6)	26–9 479–85 1618–22 708–14 795–801 1055–61 1235–45 1095–101 1490–8

Bible KC, Suman VJ, Molina JR, Smallridge RC, Maples WJ, Menefee ME, Rubin J, Sideras K, Morris JC 3rd, McIver B, Burton JK, Webster KP, Bieber C, Traynor AM, Flynn PJ, Goh BC, Tang H, Ivy SP, Erlichman	Efficacy of pazopanib in progressive, radioiodine-refractory, metastatic differentiated thyroid cancers: results of a phase 2 consortium study.	Lancet Oncol	2010	11(10)	962–72
Abraham P, Acharya Sugitani I, Toda K, Yamada K, Yamamoto N, Ikenaga Listewnik MH, Birkenfeld B, Chosia M, Elbl B, Niedzialkowska K, Sawrymowicz Reverter JL, Colome E, Halperin I, Julian T, Diaz G, Mora M, Sanmarti A, Puig— Domingo Basbug M, Ozgun MT, Murat N, Batukan C, Ozcelik B, Kurtoglu Mulazimoglu M, Edis N, Tamam MO, Uyanik E, Ozpacaci Baranauskas Z, Valuckas KP, Tiskevicius Kushchayev S, Kushchayeva Y, Theodore N, Preul MC, Clark	Current and emerging treatment options for Graves' Three distinctly different kinds of papillary thyroid microcarcinoma should be The occurrence of malignant thyroid lesions in patients after radioiodine treatment due to benign thyroid diseases.	Ther Clin Risk Manag World J Surg Endokrynol Pol	2010	6	29–40
	[Comparative study of historical series of differentiated thyroid carcinoma in two tertiary hospitals in Spain versus North American series].	Endocrinol Nutr	2010	57(8)	364–9
	Prenatal diagnosis of fetal hypothyroidism after maternal radioactive iodine exposure	J Clin Ultrasound	2010	38(9)	506–8
Mulazimoglu M, Edis N, Tamam MO, Uyanik E, Ozpacaci Baranauskas Z, Valuckas KP, Tiskevicius Kushchayev S, Kushchayeva Y, Theodore N, Preul MC, Clark	The evaluation of the external dose measurement of the patients treated with [Outcomes of long-term combined treatment in follicular thyroid carcinoma].	Radiat Prot Dosimetry	2010	141(3)	233–8
Silva F, Laguna R, Nieves-Rivera Baroli A, Pedrazzini L, Lomuscio G, Marzoli So YK, Son YI, Hong SD, Seo MY, Baek CH, Jeong HS, Chung Gerrard GE, O'Toole L, Roberts Bertagna F, Giubbini Malterling RR, Andersson RE, Falkmer S, Falkmer U, Nilehn E, Jarhult	Percutaneous vertebroplasty for thyroid cancer metastases to the spine.	Medicina (Kaunas)	2010	46(4)	268–74
	Pediatric thyroid cancer with extensive disease in a Hispanic Anaplastic thyroid carcinoma.	Thyroid	2010	20(5)	555–60
	Practical aspects of multimodal therapy and data emerging from Subclinical lymph node metastasis in papillary thyroid microcarcinoma: a study of 551	J Pediatr Endocrinol Minerva Endocrinol	2010	23(1–2)	59–64
	Should we routinely offer a second admission for radioiodine to patients with To the editor: Long-term follow-up of patients with	Surgery Clin Oncol (R Coll Radiol) World J Surg	2010	148(3)	526–31
	Differentiated thyroid cancer in a Swedish county—long-term results and quality of life.	Acta Oncol	2010	49(4)	454–9

Verburg FA, Stokkel MP, Duren C, Verkooijen RB, Mader U, van Isselt JW, Marlowe RJ, Smit JW, Reiners C, Grant CS, Stulak JM, Thompson GB, Richards ML, Reading CC, Hay Yipintsoi T, Premprabha T, Geater A, Thientunyakij T, Asli IN, Baharfard N, Shafiei B, Tabei F, Javadi H, Seyedabadi M, Nabipour I, Assadi Wang J, Yuan H, Ma Q, Liu X, Wang H, Jiang Y, Tian S, Yang Tuttle RM, Rondeau G, Lee Verburg FA, Luster M, Lassmann M, Reiners Hurtado-Lopez LM, Melchor-Ruan J, Basurto-Kuba E, Montes de Oca-Duran ER, Pulido-Cejudo A, Athie-Fallahi B, Adabi K, Majidi M, Fard-Esfahani A, Heshmat R, Larjani B, Haghpanah Lubin Papendieck P, Gruneiro-Papendieck L, Venara M, Acha O, Maglio S, Bergada I, Chiesa Ozkan E, Soydal C, Araz M, Kucuk Nixon IJ, Ganly I, Patel S, Palmer FL, Whitcher MM, Tuttle RM, Shah Vaisman F, Tala H, Grewal R, Tuttle	No survival difference after successful (131)I ablation between patients with initially low-risk and high-risk differentiated thyroid cancer. Risks and adequacy of an optimized surgical approach to the primary surgical management of papillary thyroid Mortality-related factors in 1056 radioiodine-treated patients with well-differentiated thyroid cancer in southern Thailand. Relation between clinical and laboratory parameters with radiation dose rates from patients receiving iodine-131 Interstitial 125I seeds implantation to treat spinal metastatic and primary tumor. A risk-adapted approach to the use of radioactive iodine and external beam radiation in the (131)I therapy in patients with benign thyroid disease does not conclusively lead to a higher Low-risk papillary thyroid cancer recurrence in patients treated with total thyroidectomy and adjuvant therapy vs. patients treated with partial thyroidectomy. Incidence of second primary malignancies during a long-term surveillance of patients with differentiated thyroid carcinoma in relation to radioiodine Radioactive iodine 1311 (RAI) treatment. The nearest to the "magic bullet" but should Differentiated thyroid carcinoma: presentation and follow-up in children and adolescents. Differentiated thyroid carcinomas in childhood: The impact of microscopic extrathyroid extension on outcome in patients with clinical T1 and T2 well-differentiated thyroid cancer, an incomplete structural response to therapy is associated with significantly	Eur J Nucl Med Mol Imaging World J Surg World J Surg Radiat Prot Dosimetry Med Oncol Cancer Control Nuklearmedizin Cir Cir Clin Nucl Med Pediatr Endocrinol Rev J Pediatr Endocrinol Metab J Pediatr Endocrinol Surgery Thyroid	2010 2010 2010 2010 2010 2011 2011 2011 2011 2011 2011 2011 2011 2011	37(2) 34(6) 34(2) 138(4) 27(2) 18(2) 50(3) 79(2) 36(4) 9(1) 24(9–10) 24(9–10) 150(6) 21(12)	276–83 1239–46 230–6 376–81 319–26 89–95 93–9; quiz N20 118–25 277–82 415–6 743–8 739–42 1242–9 1317–22
--	--	--	--	--	--

Soyluk O, Boztepe H, Aral F, Alagol F, Ozbeyp	Papillary thyroid carcinoma patients assessed to be at low or intermediary risk after primary treatment are at greater risk of long term	Thyroid	2011	21(12)	1301–8
Kim EY, Kim TY, Kim WG, Yim JH, Han JM, Ryu JS, Hong SJ, Yoon JH, Gong G, Kim WB, Shong Castagna MG, Maino F, Cipri C, Belardini V, Theodoropoulou A, Cevenini G, Pacini Huang BY, Lin JD, Chao TC, Lin KJ, Hseuh C, Tsang Iyer NG, Morris LG, Tuttle RM, Shaha AR, Ganly van Dijk D, Plukker JT, van der Horst-Schrivers AN, Jansen L, Brouwers AH, Muller-Kobold A, Sluiter WJ, Links Hieu TT, Russell AW, Cuneo R, Clark J, Kron T, Hall P, Doi Haugen Zhao Y, Zhang Y, Liu XJ, Shi Bal C, Chandra P, Kumar A, Dwivedi Daumerie C, Boschi A, Perros Mallick U, Harmer C, Hackshaw A, Moss Adedapo KS, Fadiji IO, Orunmuyi AT, Ejeh JE, Osifo Brownlie BJ, Turner J, Abdelaal Ibrahimasic T, Nixon IJ, Palmer FL, Whitcher MM, Tuttle RM, Shaha A, Patel SG, Shah JP, Ganly Hugo J, Robenshtok E, Grewal R, Larson S, Tuttle Lim I, Kim SK, Hwang SS, Kim SW, Chung KW, Kang HS, Lee	Effects of different doses of radioactive iodine for remnant ablation on successful ablation and on long-term recurrences in patients with differentiated Delayed risk stratification, to include the response to initial treatment (surgery and radioiodine ablation), has better Therapeutic outcomes of papillary thyroid cancer patients in different risk groups. Rising incidence of second cancers in patients with low-risk (T1N0) thyroid cancer who The value of detectable thyroglobulin in patients with differentiated thyroid cancer after initial (1)(3)(1)I therapy. Cancer risk after medical exposure to radioactive iodine in benign thyroid diseases: a Radioiodine remnant ablation: current indications and dosing Prognostic factors for differentiated thyroid carcinoma A randomized equivalence trial to determine the optimum dose of iodine-131 for remnant Is Recombinant Human TSH a Trigger for Graves' Iodine or Not (IoN) for low-risk differentiated thyroid cancer: the next UK National Cancer High default rate in thyroid cancer management in Ibadan, Nigeria: a need for health Deaths due to differentiated thyroid cancer: a South Island, Undetectable thyroglobulin after total thyroidectomy in patients with low- and intermediate-risk papillary thyroid cancer—is there a need Recombinant human thyroid stimulating hormone-assisted radioactive iodine remnant ablation in thyroid cancer Prognostic implication of thyroglobulin and quantified whole body scan after initial radioiodine therapy on early	Nucl Med Commun Eur J Endocrinol Oncology Cancer Clin Endocrinol (Oxf) Endocr Relat Cancer Endocr Pract Tumori Nucl Med Commun Eur Thyroid J Clin Oncol (R Coll Radiol) Afr J Med Med Sci N Z Med J Surgery Thyroid Ann Nucl Med	2011 2011 2011 2011 2011 2011 2011 2012 2012 2012 2012 2012 2012 2012 2012 2012 2012 2012 2012 2012 2012 2012 2012 2012	32(10) 165(3) 80(1–2) 117(19) 74(1) 19(5) 18(4) 98(2) 33(10) 1(2) 24(3) 41 Suppl 105–9 125(136) 13–21 152(6) 22(10) 26(10)	954–9 441–6 123–9 4439–46 104–10 645–55 604–10 233–7 1039–47 105–9 159–61 1007–15 777–86

Rosenbaum-Krumme SJ, Gorges R, Bockisch A, Binse Kutluhan A, Yalciner G, Bozdemir K, Ozdemir E, Tarlak B, Bilgen Ozpacaci T, Mulazimoglu M, Tamam MO, Leblebici C, Yildiz K, Uyanik E, Hartl DM, Leboulleux S, Al Ghuzlan A, Baudin E, Chami L, Schlumberger M, Travagli Kim K, Kim SJ, Kim IJ, Kim YK, Kim BS, Pak Enomoto Y, Enomoto K, Uchino S, Shibuya H, Watanabe S, Noguchi Barbesino G, Goldfarb M, Parangi S, Yang J, Ross DS, Daniels Schwartz C, Bonneta F, Dabakuyo S, Gauthier M, Cueff A, Fieffe S, Pochart JM, Cochet I, Crevisy E, Dalac A, Papathanassiou D, Toubeau Lang BH, Wong IO, Wong KP, Cowling BJ, Wan Qin C, Cau W, Zhang Y, Mghanga FP, Lan X, Gao Z, An Vrachimis A, Schober O, Riemann Huang CH, Chao TC, Hseuh C, Lin KJ, Ho TY, Lin SF, Lin de Meer SG, Dauwan M, de Keizer B, Valk GD, Borel Rinkes IH, Vriens Ito Y, Kudo T, Kobayashi K, Miya A, Ichihara K, Miyauchi Wu G, Fraser S, Pai SI, Farrag TY, Ladenson PW, Tufano	(1)(8)F-FDG PET/CT changes therapy management in high-risk DTC after first radioiodine Papillary thyroid carcinoma with metastasis to the temporooccipital skull: a case report. Intraocular and orbital metastasis as a rare form of clinical presentation of insular thyroid cancer. Optimization of staging of the neck with prophylactic central and lateral neck dissection for papillary thyroid carcinoma. Clinical significance of diffuse hepatic visualization and thyroid bed uptake on post-ablative Clinical features, treatment, and long-term outcome of papillary thyroid cancer in children and adolescents without radiation Thyroid lobe ablation with radioactive iodine as an alternative to completion thyroidectomy after Impact on overall survival of radioactive iodine in low-risk differentiated thyroid cancer patients. Risk of second primary malignancy in differentiated thyroid carcinoma treated with Correlation of clinicopathological features and expression of molecular Radioiodine remnant ablation in differentiated thyroid cancer after combined endogenous and Therapeutic outcome and prognosis in young patients with papillary and follicular thyroid Not the number but the location of lymph nodes matters for recurrence rate and disease-free survival in patients Prognostic factors for recurrence of papillary thyroid carcinoma in the lymph nodes, Determining the extent of lateral neck dissection necessary to establish regional disease control and avoid	Eur J Nucl Med Mol Imaging Kulak Burun Bogaz Ihtis Derg Ann Endocrinol (Paris) Ann Surg Onkologie World J Surg Thyroid J Clin Endocrinol Metab Surgery Clin Nucl Med Nuklearmedizin Pediatr Surg Int World J Surg World J Surg Head Neck	2012 2012	39(9) 22(3) 73(3) 255(4) 35(3) 36(6) 22(4) 97(5) 151(6) 37(3) 51(3) 28(5) 36(6) 36(6) 34(10)	1373–80 160–3 222–4 777–83 82–6 1241–6 369–76 1526–35 844–50 e40–6 67–72 489–94 1262–7 1274–8 1418–21
---	--	--	--	--	---

Vianello F, Mazzarotto R, Mian C, Lora O, Saladini G, Servadio O, Basso M, Pennelli G, Pelizzo Sun XS, Guevara N, Fakhry N, Sun SR, Marcy PY, Santini J, Bosset JF, Thariat	Clinical outcome of low-risk differentiated thyroid cancer patients after radioiodine remnant ablation and recombinant human thyroid-	Clin Oncol (R Coll Radiol)	2012	24(3)	162–8
Lang BH, Ng SH, Lau LL, Cowling BJ, Wong KP, Wan	[Radiation therapy in thyroid cancer].	Cancer Radiother	2013	17(3)	233–43; quiz 255–6, 25
Azizmohammadi Z, Tabei F, Shafiei B, Babaei AA, Jukandan SM, Naghshine R, Javadi H, Nabipour I, Assadi M, Asli	A systematic review and meta-analysis of prophylactic central neck dissection on short-term locoregional recurrence in	Thyroid	2013	23(9)	1087–98
An YS, Yoon JK, Lee SJ, Song HS, Yoon SH, Jo	A study of the time of hospital discharge of differentiated thyroid cancer patients after receiving iodine-131 for thyroid remnant ablation treatment.	Hell J Nucl Med	2013	16(2)	103–6
Kim HJ, Kim NK, Choi JH, Kim SW, Jin SM, Suh S, Bae JC, Min YK, Chung JH, Kim Kruijff S, Aniss AM, Chen P, Sidhu SB, Delbridge LW, Robinson B, Clifton- Bligh RJ, Roach P, Gill AJ, Learoyd D,	Symptomatic late-onset sialadenitis after radioiodine therapy in thyroid cancer. Radioactive iodine ablation does not prevent recurrences in patients with papillary thyroid microcarcinoma.	Ann Nucl Med	2013	27(4)	386–91
Liu WS, Zhang GF, Xu Wienhold R, Scholz M, Adler JR, G Nster C, Paschke Young S, Harari A, Smooke-Praw S, Ituarte PH, Yeh Pedrazzini L, Baroli A, Marzoli L, Guglielmi R, Papini Vrachimis A, Riemann B, Gerss J, Maier T, Schober Saengsuda	Decreasing the dose of radioiodine for remnant ablation does not increase structural recurrence rates in papillary thyroid carcinoma. [Management and prognostic factors for 119 patients with The management of thyroid nodules: a retrospective analysis of health insurance Effect of reoperation on outcomes in papillary thyroid cancer]. Cancer recurrence in papillary thyroid microcarcinoma: a multivariate analysis on 231 Peace of mind for patients with differentiated thyroid cancer? Radioiodine remnant ablation in low-risk differentiated thyroid	Clin Endocrinol (Oxf)	2013	78(4)	614–20
Park S, Jeong JS, Ryu HR, Lee CR, Park JH, Kang SW, Jeong JJ, Nam KH, Chung WY, Park	Surgery	2013	154(6)	1337–44; discussion 1;	
Zhonghua Zhong Liu	Zhonghua Zhong Liu	2013	35(10)	778–82	
Dtsch Arztebl Int	Dtsch Arztebl Int	2013	110(49)	827–34	
Surgery	Surgery	2013	154(6)	1354–61; discussion 1;	
Minerva Endocrinol	Minerva Endocrinol	2013	38(3)	269–79	
Nuklearmedi zin	Nuklearmedi zin	2013	52(4)	115–20	
J Med Assoc Thai	J Med Assoc Thai	2013	96(5)	614–24	
J Korean Med Sci	J Korean Med Sci	2013	28(5)	693–9	

Reiners C, Biko J, Haenscheid H, Hebestreit H, Kirinjuk S, Baranowski O, Marlowe RJ, Demidchik E, Drozd V, Demidchik	Twenty-five years after Chernobyl: outcome of radioiodine treatment in children and adolescents with very high-risk radiation-induced differentiated thyroid carcinoma.	J Clin Endocrinol Metab	2013	98(7)	3039–48
Kuo EJ, Goffredo P, Sosa JA, Roman	Aggressive variants of papillary thyroid microcarcinoma are associated with extrathyroidal	Thyroid	2013	23(10)	1305–11
Castagna MG, Cevenini G, Theodoropoulou A, Maino F, Memmo S, Claudia C, Belardini V, Brianzoni E, Pacini	Post-surgical thyroid ablation with low or high radioiodine activities results in similar outcomes in intermediate risk differentiated thyroid cancer patients.	Eur J Endocrinol	2013	169(1)	23–9
Meixner M, Hellmich M, Dietlein M, Kobe C, Schicha H, Schmidt	Disease-free survival in papillary and follicular thyroid carcinoma. Comparison between UICC 5th and 7th	Nuklearmedizin	2013	52(3)	71–80
Lango M, Flieder D, Arrangoiz R, Veloski C, Yu JQ, Li T, Burtness B, Mehra R, Galloway T, Ridge	Extranodal extension of metastatic papillary thyroid carcinoma: correlation with biochemical endpoints, nodal persistence, and systemic	Thyroid	2013	23(9)	1099–105
Hod R, Bachar G, Sternov Y, Shvero	Insular thyroid carcinoma: a retrospective clinicopathologic	Am J Otolaryngol	2013	34(4)	292–5
Ito Y, Hirokawa M, Masuoka H, Yabuta T, Kihara M, Higashiyama T, Takamura Y, Kobayashi K, Miya A, Ibrahimasic T,	Prognostic factors of minimally invasive follicular thyroid carcinoma: extensive vascular invasion significantly affects patient prognosis.	Endocr J	2013	60(5)	637–42
Ghossein R, Carlson DL, Chernichenko N, Nixon I, Palmer FL, Lee NY, Shaha AR, Patel SG, Tuttle RM, Balm AJ, Shah JP, Ganly	Poorly differentiated thyroid carcinoma presenting with gross extrathyroidal extension: 1986–2009 Memorial Sloan-Kettering Cancer Center experience.	Thyroid	2013	23(8)	997–1002
Amdur RJ, Dan T, Mazzaferri	Absence of bone marrow toxicity in elderly patients treated with recombinant human thyroid-stimulating	Am J Clin Oncol	2013	36(4)	348–53
Hilly O, Stern-Shavit S, Iran S, Feinmesser	Treatment decisions and adherence to guidelines in the treatment of low risk papillary	Isr Med Assoc J	2014	16(9)	548–52
Rokni H, Sadeghi R, Moossavi Z, Treglia G, Zakavi	Efficacy of different protocols of radioiodine therapy for treatment of toxic nodular	Int J Endocrinol Metab	2014	12(2)	e14424
Wang TS, Goffredo P, Sosa JA, Roman Iacobone M, Jansson S, Barczynski M, Goretzki	Papillary thyroid microcarcinoma: an over-Multifocal papillary thyroid carcinoma—a consensus report of the European Society of	World J Surg	2014	38(9)	2297–303
Karyampudi A, Hamide A, Halanaik D, Sahoo JP,	Radioiodine therapy in patients with Graves' disease and the effects of prior carbimazole	Langenbecks Arch Surg	2014	399(2)	141–54
		Indian J Endocrinol Metab	2014	18(5)	688–93

Verburg FA, Mader U, Reiners C, Hanscheid	Long-term survival in differentiated thyroid cancer is worse after low-activity initial	J Clin Endocrinol Metab	2014	99(12)	4487–96
Guy A, Hirsch D, Shohat T, Bachar G, Tirosh A, Robenshtok E, Shimon I, Benbassat	Papillary thyroid cancer: factors involved in restaging N1 disease after total thyroidectomy and radioactive iodine treatment.	J Clin Endocrinol Metab	2014	99(11)	4167–73
Liu B, Peng W, Huang R, Tian R, Zeng Y, Kuang	Thyroid cancer: radiation safety precautions in ^{131}I therapy based on actual biokinetic	Radiology	2014	273(1)	211–9
Ito Y, Miyauchi A, Ito M, Yabuta T, Masuoka H, Higashiyama T, Fukushima M, Kobayashi K, Kihara Lupoli R, Cacciapuoti M, Tortora A, Barba L, Verde N, Romano F, Vastarella M, Fonderico F, Masone S, Milone M, Lupoli G, Lupoli Tresoldi AS, Sburlati LF, Rodari M, Schinkelshoek M, Perrino M, De Leo S, Montefusco L, Colombo P, Arosio M, Lania AG, Fugazzola L, Chiti	Prognosis and prognostic factors of differentiated thyroid carcinoma after the appearance of metastasis refractory to radioactive iodine therapy.	Endocr J	2014	61(8)	821–4
Astl J, Chovanec M, Lukes P, Katra R, Dvorakova M, Vlcek P, Sykorova P, Betka Hasbek Z, Turgut B, Kilicli F, Altuntas EE, Yucel Ardito G, Avenia N, Giustozzi E, Salvatori M, Fadda G, Ardito F, Revelli De Crea C, Raffaelli M, Sessa L, Ronti S, Fadda G, Bellantone C, Lombardi	Clinical outcome in differentiated thyroid carcinoma Int J Surg and microcarcinoma.	Int J Pediatr Otorhinolaryngol	2014	37(8)	709–14
Lee J, Song Y, Soh Han JM, Kim WG, Kim TY, Jeon MJ, Ryu JS, Song DE, Hong SJ, Shong YK, Kim Goldfarb M, Sener Rosario PW, Calsolari	Thyroid carcinoma surgery in children and adolescents – 15 years experience surgery of pediatric thyroid carcinoma. Importance of postoperative stimulated thyroglobulin level at the time of ^{131}I ablation Papillary thyroid microcarcinoma: proposal of treatment based on histological prognostic factors evaluation. Actual incidence and clinical behaviour of follicular thyroid carcinoma: an institutional experience. Central lymph node metastasis is an important prognostic factor in patients with papillary thyroid carcinoma Effects of low-dose and high-dose postoperative radioiodine therapy on the clinical outcome in patients with small Comparison of radioiodine utilization in adolescent and Thyroid ablation with 1.1 GBq (30 mCi) iodine-131 in patients with papillary thyroid carcinoma	Asian Pac J Cancer Prev	2014	15(6)	2523–7
		Ann Ital Chir	2014	85(1)	1–5
		ScientificWorldJournal	2014	2014	952095
		J Korean Med Sci	2014	29(1)	48–52
		Thyroid	2014	24(5)	820–5
		Endocr Pract	2014	20(5)	405–11
		Thyroid	2014	24(5)	826–31

Kotb MH, Zaher AM, Abd El-Wahab MA, Abulkheir IL, Hussein M, Salem	Prognostic value of p27 in follicular thyroid carcinoma and its relation to radioactive iodine response: does it aid in the	Appl Immunohist ochem Mol Morphol	2014	22(7)	511–7
Kim SK, Yun GY, Kim KH, Park SK, Choi HY, Ha SK, Park Jeon MJ, Kim WG, Park WR, Han JM, Kim TY, Song DE, Chung KW, Ryu JS, Hong SJ, Shong YK, Zoghlami A, Roussel F, Sabourin JC, Kuhn JM, Marie JP, Dehesdin D, Choussy	Severe hyponatremia following radioactive iodine therapy in patients with differentiated Modified dynamic risk stratification for predicting recurrence using the response to initial therapy in patients with differentiated thyroid BRAF mutation in papillary thyroid carcinoma: predictive value for long-term prognosis and radioiodine sensitivity. Prognostic factors affecting	Thyroid	2014	24(4)	773–7
Akkas BE, Demirel BB, Vural	disease-specific survival in patients with recurrent and/or metastatic differentiated	Eur J Endocrinol	2014	170(1)	23–30
Biondi B, Bartalena L, Cooper DS, Hegedus L, Laurberg P, Kahaly	The 2015 European Thyroid Association Guidelines on Diagnosis and Treatment of	Eur Ann Otorhinolary ngol Head Neck Dis	2014	131(1)	7–13
Xu G, Wu T, Ge L, Li Ruel E, Thomas S, Dinan M, Perkins JM, Roman SA, Sosa	A Systematic Review of Adjuvant Interventions for Adjuvant radioactive iodine therapy is associated with improved survival for patients	Oncol Res Treat	2015	38(7–8)	368–72
Sacks W, Wong RM, Bresee C, Braunstein	Use of evidence-based guidelines reduces radioactive iodine treatment in patients	J Clin Endocrinol Metab	2015	100(4)	1529–36
Fu H, Ma C, Tang L, Wu F, Liu B, Wang	Recombinant human thyrotropin versus thyroid hormone withdrawal in radioiodine	Thyroid	2015	25(4)	377–85
Worden F, Fassnacht M, Shi Y, Hadjiev T, Bonichon F, Gao M, Fugazzola L, Ando Y, Hasegawa Y, Park DJ, Shong YK, Smit JW, Chung J, Kappeler C, Meinhardt G, Schlumberger M, Sawka AM, Straus S, Rodin G, Thorpe KE, Ezzat S, Gafni A, Goldstein	Safety and tolerability of sorafenib in patients with radioiodine-refractory thyroid cancer.	Q J Nucl Med Mol Imaging	2015	59(1)	121–8
Jacob JJ, Stephen C, Paul TV, Thomas N, Oommen R, Seshadri Czepczynski R, Matysiak-Grzes M, Gryczynska M, Baczyk M, Wyszomirska A, Shen J, Wang S, Zhao X, Shao X, Jiang X, Dai Y, Xu S	Decision aid on radioactive iodine treatment for early stage papillary thyroid cancer: update to study protocol with follow- No impact of dietary iodine restriction in short term development of hypothyroidism	Endocr Relat Cancer	2015	22(6)	877–87
	Peptide receptor radionuclide therapy of differentiated thyroid cancer: efficacy and toxicity.	Trials	2015	16	302
	Skull metastasis from follicular thyroid carcinoma: report of three cases and review of	Indian J Endocrinol Metab	2015	19(1)	60–5
		Arch Immunol Ther Exp (Warsz)	2015	63(2)	147–54
		Int J Clin Exp Pathol	2015	8(11)	15285–93

Kang	Using ultrasound radio frequency technology to assess regression of the structure and function of the carotid artery	Arch Med Sci	2015	11(6)	1236–43
Kao YH, Gan HK, Zaheer S, Lam WW, Loke KS, Wong WY, Ng DC, Goh	Gender, Race, and Age at Diagnosis as Risk Factors for Metastasis or Recurrence among 1,657 Thyroid Cancer	Oncol Res Treat	2015	38(12)	679–82
Al-Qahtani KH, Al Asiri M, Tunio MA, Aljohani NJ, Bayoumi Y, Fatani H, AlHadab Z, Zaman MU, Fatima N, Zaman U, Sajjad Z, Zaman A, Tahseen	Adjuvant Radioactive iodine 131 J ablation in papillary microcarcinoma of thyroid: Saudi Arabian experience Predictive value of pyramidal lobe, percentage thyroid uptake and age for ablation outcome SERIAL NECK ULTRASOUND	Otolaryngol Head Neck Surg Indian J Nucl Med	2015	44	51
Yang SP, Bach AM, Tuttle RM, Fish	IS MORE LIKELY TO IDENTIFY FALSE-POSITIVE ABNORMALITIES THAN	Endocr Pract	2015	21(12)	1372–9
Ryödi E, Metso S, Jaatinen P, Huhtala H, Saaristo R, Valimaki M, Auvinen Liu FH, Kuo SF, Hsueh C, Chao TC, Lin	Cancer Incidence and Mortality in Patients Treated Either With RAI or Thyroidectomy for Hyperthyroidism. Postoperative recurrence of papillary thyroid carcinoma with lymph node metastasis.	J Clin Endocrinol Metab J Surg Oncol	2015	100(10)	3710–7
Ahn HY, Min HS, Yeo Y, Ma SH, Hwang Y, An JH, Choi HS, Keam B, Im SA, Park DJ, Park IA, Noh DY, Youn YK, Chung JK, Cho BY, Park SK, Seo GH, Cho YY, Chung JH, Kim	Radioactive Iodine Therapy Did Not Significantly Increase the Incidence and Recurrence of Subsequent Breast Cancer. Increased Risk of Leukemia After Radioactive Iodine Therapy in Patients with Increased risk of second primary malignancy in pediatric and young adult patients	J Clin Endocrinol Metab Thyroid	2015	100(9)	3486–93
Marti JL, Jain KS, Morris	Post-operative stimulated thyroglobulin and neck ultrasound as personalized criteria for risk stratification	Thyroid	2015	25(6)	681–7
Orlov S, Salari F, Kashat L, Freeman JL, Vescan A, Witterick IJ, Walfish Gamper EM, Wintner LM, Rodrigues M, Buxbaum S, Nilica B, Singer S, Giesinger JM, Holzner B, Virgolini	Persistent quality of life impairments in differentiated thyroid cancer patients: results from a monitoring programme.	Endocrine Eur J Nucl Med Mol Imaging	2015	50(1)	130–7
Semrad TJ, Semrad AM, Farwell DG, Chen Y, Cress	Initial treatment patterns in younger adult patients with differentiated thyroid cancer in	Thyroid	2015	25(5)	509–13
Avram AM, Esfandiari NH, Wong	Preablation 131-I scans with SPECT/CT contribute to thyroid cancer risk stratification	J Clin Endocrinol Metab	2015	100(5)	1895–902
Lorenz R, Buck A, Reiners	[In-patient nuclear medicine therapy in Germany from 2010	Nuklearmedizin	2015	54(2)	61–8

Kiernan CM, Parikh AA, Parks LL, Solorzano	Use of radioiodine after thyroid lobectomy in patients with differentiated thyroid cancer: Lateral Neck Lymph Node	J Am Coll Surg	2015	220(4)	617–25
Wang LY, Palmer FL, Nixon IJ, Tuttle RM, Shah JP, Patel SG, Shaha AR, Ganly	Characteristics Prognostic of Outcome in Patients with Clinically Evident N1b Papillary	Ann Surg Oncol	2015	22(11)	3530–6
Maier TM, Schober O, Gerss J, Gorlich D, Wenning C, Schaefers M, Riemann B,	Differentiated thyroid cancer patients more than 60 years old paradoxically show an increased life expectancy.	J Nucl Med	2015	56(2)	190–5
Shivaprasad C, Prasanna Kumar	Long-term carbimazole pretreatment reduces the Clinical management and outcomes in patients with hyperfunctioning distant metastases from differentiated	Indian J Endocrinol	2015	19(1)	84–8
Qiu ZL, Shen CT, Luo	Differentiated thyroid cancer in patients over 60 years of age at presentation: a retrospective	Thyroid	2015	25(2)	229–37
Garg A, Chopra S, Ballal S, Soundararajan R, Bal Khang AR, Cho SW, Choi HS, Ahn HY, Yoo WS, Kim KW, Kang KW, Yi KH, Park DJ, Lee DS, Chung JK, Cho BY, Park	The risk of second primary malignancy is increased in differentiated thyroid cancer patients with a cumulative (131)I dose over 37 GBq.	J Geriatr Oncol	2015	6(1)	29–37
Kim KJ, Kim SM, Lee YS, Chung WY, Chang HS, Park	Prognostic significance of tumor multifocality in papillary thyroid carcinoma and its relationship with primary tumor	Clin Endocrinol (Oxf)	2015	83(1)	117–23
Su DH, Chang SH, Chang	The impact of locoregional recurrences and distant metastases on the survival of	Ann Surg Oncol	2015	22(1)	125–31
Scott E, Learoyd D, Clifton-Bligh	Therapeutic options in papillary thyroid carcinoma: current [Ablative Radioiodotherapy bei niedrigem und intermediarem Rezidivrisiko. Hohere	Future Oncol	2016	12(22)	2603–2613
Dietlein M, Drzezga	Radiiodine therapy versus antithyroid drugs in Graves' disease: a meta-analysis of RISK FACTORS FOR	Nuklearmedizin	2016	55(3)	71–6
Wang J, Qin	NONREMISSION AND PROGRESSION-FREE SURVIVAL AFTER I-131 THERAPY IN PATIENTS WITH LUNG METASTASIS FROM	Br J Radiol	2016		20160418
Chen P, Feng HJ, Ouyang W, Wu JQ, Wang J, Sun YG, Xian JL, Huang	Risk of second primary breast cancer after radioactive iodine treatment in thyroid cancer: a	Endocr Pract	2016	22(9)	1048–56
Zhang Y, Liang J, Li H, Cong H, Lin	Low iodine diet in differentiated thyroid cancer: a review. [Lenvatinib in radioiodine refractory thyroid carcinomas].	Nucl Med Commun	2016	37(2)	110–5
Li JH, He ZH, Bansal V, Hennessey		Clin Endocrinol	2016	84(1)	3–12
de la Fouchardiere		Bull Cancer	2016	103(11)	905–910

Klein Hesselink EN, Brouwers AH, de Jong JR, van der Horst-Schrivers AN, Coppes RP, Lefrandt JD, Jager PL, Vissink Gao X, Zhang X, Zhang Y, Hua W, Maimaiti Y, Gao Brose MS, Cabanillas ME, Cohen EE, Wirth LJ, Riehl T, Yue H, Sherman SI, Sherman Tun NN, Beckett G, Zammitt NN, Strachan MW, Seckl la Cour JL, Andersen UB, Sorensen CH, Nygaard B, Jensen Canto AU, Dominguez PN, Jimeno CA, Obaldo Louvet C, De Bellis A, Pereira B, Bournaud C, Kelly A, Maqdasy S, Roche B, Desbiez F, Borson-Chazot F, Tauveron I, Batisse-Lignier Hollingsworth B, Senter L, Zhang X, Brock GN, Jarjour W, Nagy R, Brock P, Coombes KR, Kloos RT, Ringel MD, Sipos J, Lattimer I, Carrau R, Jhiang Deutschmann MW, Chin-Lenn L, Nakoneshny SC, Dort JC, Pasieka JL, Chandarana Mohan V, Lind Matthews TJ, Chua E, Gargya A, Clark J, Gao K, Elliott Banerjee M, Wiebel JL, Guo C, Gay B, Haymart	Effects of Radioiodine Treatment on Salivary Gland Function in Patients with Differentiated Thyroid Carcinoma: A Prospective Study. Is papillary thyroid microcarcinoma an indolent tumor?: A retrospective study Vemurafenib in patients with BRAF(V600E)-positive metastatic or unresectable papillary thyroid cancer Thyrotropin Receptor Antibody Levels at Diagnosis and After Thionamide Course Predict Radioiodine Therapy Does Not Change the Atherosclerotic Burden of the Carotid Arteries. Comparison of Fixed versus Calculated Activity of Radioiodine for the Treatment Time course of Graves' orbitopathy after total thyroidectomy and radioiodine therapy for thyroid cancer. Risk Factors of ^{131}I -Induced Salivary Gland Damage in Thyroid Cancer Patients. Practice patterns among thyroid cancer surgeons: implications of performing a prophylactic central neck dissection. A review of treatment options for Graves' disease: why total thyroidectomy is a viable option Elevated serum thyroglobulin levels at the time of ablative radioactive iodine therapy indicate a worse prognosis in Use of imaging tests after primary treatment of thyroid cancer in the United States: population based retrospective	J Nucl Med Medicine (Baltimore) Lancet Oncol Thyroid Thyroid Endocrinol Metab (Seoul) Medicine (Baltimore) J Clin Endocrinol Metab J Otolaryngol Head Neck Surg J Community Hosp Intern J Laryngol Otol BMJ	2016 57(11) 1685–1691 2016 95(40) e5067 2016 17(9) 1272–82 2016 26(8) 1004–9 2016 26(7) 965–71 2016 31(1) 168–73 2016 95(48) e5474 2016 101(11) 4085–4093 2016 45(1) 55 2016 6(4) 32369 2016 130 Supl S50–3 2016 354 i3839
--	---	--	---

Cramon P, Winther KH, Watt T, Bonnema SJ, Bjorner JB, Ekholm O, Groenvold M, Hegedus L, Feldt-Rasmussen U, Rasmussen	Quality-of-Life Impairments Persist Six Months After Treatment of Graves' Hyperthyroidism and Toxic Nodular Goiter: A Prospective Cohort Study.	Thyroid	2016	26(8)	1010–8
Rosario PW, Mourao GF, Calsolari	Efficacy of adjuvant therapy with 3.7 GBq radioactive iodine in intermediate-risk patients with 'higher risk features' and Patients with high-risk differentiated thyroid cancer have a lower I-131 ablation success rate than low-risk ones in spite of a high ablation	Nucl Med Commun	2016	37(11)	1148–53
Winter J, Winter M, Krohn T, Heinzel A, Behrendt FF, Tuttle RM, Mottaghy FM, Verburg Klein Hesselink MS, Nies M, Bocca G, Brouwers AH, Burgerhof JG, van Dam EW, Havekes B, van den Heuvel-Eibrink MM, Corssmit EP, Kremer LC, Netea-Maier RT, van der Pal HJ, Peeters RP, Schmid KW, Smit JW, Williams GR, Plukker JT, Ronckers CM, van Santen HM, Tissing WJ, Links Lin CM, Yeh PT, Doyle P, Tsan YT, Chen Ruhlmann M, Binse I, Bockisch A, Rosenbaum-Krumme Heaton CM, Chang JL, Orloff Rosario PW, Mourao GF, Calsolari Prpic M, Kruljac I, Kust D, Kirigin LS, Jukic T, Dabelic N, Bolanca A, Kusic Lin CY, Lin CL, Huang WS, Kao Pathak KA, Klonisch TC, Nason So K, Smith RE, Davis Lee DY, Won JK, Lee SH, Park DJ, Jung KC, Sung MW, Wu HG, Kim KH, Park YJ	Pediatric Differentiated Thyroid Carcinoma in The Netherlands: A Nationwide Follow-Up Study. Association Between 131I Treatment for Thyroid Cancer and Risk of Receiving Cataract Initial [18F]FDG PET/CT in high-risk DTC patients. A three-year follow-up. Prognostic Implications of Lymph Node Yield in Central and Lateral Neck Dissections Low postoperative nonstimulated thyroglobulin as a criterion for the indication of low radioiodine activity in Re-ablation I-131 activity does not predict treatment success in low- and intermediate-risk patients with differentiated Risk of Breast Cancer in Patients with Thyroid Cancer Receiving or Not Receiving Stage II differentiated thyroid cancer: A mixed bag. Radiotherapy in well-differentiated thyroid cancer: is Changes of Clinicopathologic Characteristics and Survival Outcomes of Anaplastic and Poorly Differentiated Thyroid	J Clin Endocrinol Metab J Nucl Med Nuklearmedizin Thyroid Clin Endocrinol (Oxf) J Clin Endocrinol J Nucl Med Endocrine J Surg Oncol ANZ J Surg Thyroid	2016 2016 2016 2016 2016 2016 2016 2016 2016 2016 2016 2016 2016 2016 2016 2016 2016 2016 2016	101(5) 37(11) 85(6) Metab 101(5) 57(6) 55(3) 26(3) 85(3) 52(3) 57(5) 113(1) 86(9) 26(3)	926–931 1148–53 2031–9 836–41 99–103 434–40 453–8 602–8 685–90 94–7 696–700 404–13

Vagney M, Desquilibet L, Reyes-Gomez E, Delisle F, Devauchelle P, Rodriguez-Pineiro MI, Rosenberg D, de Al Eyadah AA, Al-Sarihin KM, Etewi SM, Al-Omari AA, Al-Asa'd RA, Haddad Giesecke P, Rosenqvist M, Frykman V, Friberg L, Wallin G, Hoijer J, Lonn S, Torring Zhang H, Cai Y, Zheng L, Zhang Z, Jiang Scerrino G, Melfa G, Raspanti C, Attard A, Mazzola S, Gullo R, Bonventre S, Attard M, Cocorullo G, Gulotta Cebulska-Wasilewska A, Krzysiek M, Krajewska G, Stepien A, Krajewski Al-Qurayshi Z, Shama MA, Randolph GW, Kandil Lucy JM, Peterson ME, Randolph JF, Scrivani PV, Rishniw M, Davignon DL, Thompson MS, Scarlett Kim J, Cho SG, Kang SR, Kwon SY, Cho DH, Cho JS, Song Jeong JH, Kong EJ, Jeong SY, Lee SW, Cho IH, Ah Chun K, Lee J, Ahn Deandreas D, Rubino C, Tala H, Leboulleux S, Terroir M, Baudin E, Larson S, Fagin JA, Schlumberger M, Tuttle Parameswaran R, Shulin Hu J, Min En N, Tan WB, Yuan Sabet A, Binse I, Dogan S, Koch A, Rosenbaum-Krumme SJ, Biersack HJ, Biermann K, Ezziddin	Survival times for cats with hyperthyroidism treated with a 3.35 mCi iodine-131 dose: a retrospective study of 96 cases. Thyroid cancer post radioactive iodine treatment for hyperthyroidism (case series and review of the literature). Increased Cardiovascular Mortality and Morbidity in Patients Treated for Toxic Nodular Goiter Compared to Graves' Disease and Nontoxic Postoperative radioactive iodine-131 ablation is not necessary among patients with intermediate-risk differentiated The prevalence of post-thyroidectomy chronic asthenia: a prospective cohort study. Retrospective Biological Dosimetry at Low and High Doses of Radiation and Radioiodine Impact on Individual Minimal extrathyroidal extension does not affect survival of well-differentiated Efficacy of Low-dose (2 millicurie) versus Standard-dose (4 millicurie) Radioiodine Treatment for Cats with Mild-to-Moderate Hyperthyroidism. Preparation for radioactive iodine therapy is not a risk factor for the development of Clinical outcomes of low-dose and high-dose postoperative radioiodine therapy in patients with intermediate-risk Comparison of Empiric Versus Whole-Body/-Blood Clearance Dosimetry-Based Approach to Radioactive Iodine Treatment in Patients with Metastases from Differentiated Thyroid Cancer. Patterns of metastasis in follicular thyroid carcinoma and the difference between early Distinguishing synchronous from metachronous manifestation of distant metastases: a prognostic feature in differentiated thyroid	J Feline Med Surg Endokrynol Pol Thyroid Hell J Nucl Med Langenbecks Arch Surg Genome Integr Endocr Relat Cancer J Vet Intern Med Medicine (Baltimore) Nucl Med Commun J Nucl Med Ann R Coll Surg Engl Eur J Nucl Med Mol Imaging	2017 2017 2017 2017 2017 2017 2017 2017 2017 2017 2017 2017 2017 2017 2017 2017 2017 2017 2017 2017	2017 2017 2017 2017 2017 2017 2017 2017 2017 2017 2017 2017 2017 2017 2017 2017 2017 2017 2017 2017	1098612X17718416 878–885 3–10 1095–1102 2 221–226 326–334 e6004 228–233 717–722 151–154 190–195
--	--	---	--	--	--

5

2040

47-9

38

344-5

361-2